

HISTORICAL OVERVIEW OF THE STUDIES OF ALIEN FLORA OF UKRAINE

SHEVERA V. Myroslav^{1, 2*}, PROTOPOPOVA V. Vira^{2, 1}, BURDA I. Raisa³,
ZAVIALOVA V. Liudmyla¹, KUCHER O. Oksana¹, KORNIYENKO M. Olga¹

Abstract: This review has shown that the alien fraction of the Ukrainian flora have been identified and studied comprehensively. Thus, a total richness of alien fraction is about 1,000 species. The structure, typical trends, and regional peculiarities of this component of the Ukrainian flora have been analyzed. The main results of investigations are summarized. The alien floras of some regions, cities and towns, protected areas of Ukraine have been studied. The species composition of some habitats, partial floras with a high abundance of non-native plants, tolerance to anthropogenic influences, the segetal potential, and the role in developing of the human-made biotopes and in evolution of the plant cover have been determined. The changes in the species composition over 160 years, and the trends forming the stable and unstable elements of the alien flora of Ukraine, have been shown. The peculiarities of the distribution of archaeophytes and neophytes of the regional floras and urban floras as latitudinal gradients have been revealed. The plant invasions and expansions of 30 species have been reconstructed, analyzed and reviewed. A group of highly invasive potential species is selected. Some important characteristics of alien species, such as their activity in plant communities and their ecological effects, the trends of distribution and abundance, the participation and role under the influence of anthropogenic environmental change or climate change in various types of biotopes and geographical zones etc., have been analyzed.

Key words: Ukraine, alien flora, review, trends of investigation, peculiarities, invasive alien species

Received 09 November 2018

Accepted 17 November 2018

Introduction

The problem of alien organisms, their impact on the flora, fauna and human society attract special attention from botanists in Ukraine (Protopopova et al. 2002a), because there is more than 80% of human-made plant cover in its territory.

Complex historical factors (the invasion and relocation of nomads, the wars, the early occurrence of agriculture in the I-II centuries B.C., the settled population, the passing of important trade routes through the territory, the foundation of colonies by

¹ M.G. Kholodny Institute of Botany, NAS of Ukraine, Tereshchenkivska St., 2, 01004- Kyiv, Ukraine.

² Ferenc Rákóczi II Transcarpathian Hungarian College of Higher Education, Koshut sq., 6, 90200-Berehove, Ukraine.

³ Institute for Evolutionary Ecology, NAS of Ukraine, Lebedeva St., 2, 03143-Kyiv, Ukraine.

* Correspondence: shevera.myroslav@ukr.net

immigrants from the Mediterranean, Southern and Central Europe etc.) and further intensive development of agriculture, industry, transport, socio-economic relations, urbanization, favorable natural and climatic conditions, etc. gave wide opportunities for the invasion of alien plants (Protopopova 1973). The current state of the flora of Ukraine is characterized by a significant increase of the anthropophilic component's role in it (Protopopova et al. 2002a), which is over 15.2% of the total number of species (Fig. 1.).




Fig. 1. The increase of the number of species of alien plants in the flora of Ukraine during a period of more than 160 years

Ukraine is situated in several natural climatic zones (forest, forest-steppe, steppe); the presence of the Carpathian and the Crimean mountain ranges, a variety of natural conditions, specificity of the historical development of the country and the current ecological situation allow us to consider it as a model object for the study of the peculiarities of alien plant migration.

The process of flora adventization in different regions of the country occurs with varying intensity. The negative impact of invasive species on biodiversity is most noticeable in the regions with fragmented natural vegetation cover. The total cumulative influence of alien plants increases the insularization of populations of native plant species caused by human activity, accelerating this process significantly. There is also destructive impact to the populations of certain species or ecosystems by phytocoenically active invasive plants, established in the natural plant communities that were weakened by the anthropogenic press (Protopopova et al. 2002a, 2003). In this connection, phytinvasions continue to attract the attention of many florists; various aspects of research in this direction are developing.

To find out the current state of the alien fraction of the flora of Ukraine at the regional level, a large amount of diverse factual material was accumulated, it is characterized by the multidimensional nature of its research. This material allows us to

perform a retrospective analysis of the alien flora formation and identify the directions and trends of the further course of adventization of the plant cover of Ukraine.

Material and methods

The object of the research was the alien flora of Ukraine, and, in particular, the alien plant species which are characterized by a high degree of naturalization and discovered to have high biogeographic and coenotic activity. The main materials for the analysis and generalizations were literary sources that depicted the results of the study of alien plant species of the flora of Ukraine. The collections of herbaria have been critically studied (CHER, CWU, DNZ, DSU, KHEM, KHER, KW, KWHA, KWHU, LW, LWS, LWKS, MSUD, SIMF, UU, YALT, etc.), as well as data from the field research of the authors since 1965. The classic comparative morphological-ecological-geographical method is used in this work.

Results and discussion

The high degree of adventization of the flora of Ukraine has defined a wide range of studies. Publications on these themes from 1787 to 2017 are summarized in five issues of bibliographic references (Burda et al. 2018), which included 1697 works, of which 1183 are directly related to alien plants, and 514 – contain more or less complete information about them. The main directions of research are: floristics – 719 publications, chorology – 139, biology (primarily of weeds) – 117, ecology – 79, distribution control and control means – 75, taxonomy – 28 population – 17, general issues – 14, molecular genetics – 2, economic evaluation – 2, paleobotany – 2. The most complete studies were made on the species of the listed genera: *Ambrosia* L. – 32 publications, *Impatiens* L. – 18, *Heracleum* L. – 17, *Oenothera* L. – 14, *Phalacrolooma* Cass. – 13, *Amaranthus* L., *Solidago* L. – 11, *Asclepias* L. – 10.

The study of the alien fraction has over 220 years of history and is connected with the general research of the flora of Ukraine or its regions. Accumulation of information on alien plants began in the process of studying regional flora back in the 18th century. Initial and fragmented information about alien plants is recorded in the works of naturalists of the 18th- 19th centuries (Güldenstädt 1787, 1791, Bieberstein 1808a, b, c, Besser 1809, 1822, Andrzejowski 1823, 1860, 1862, Zawadzki 1835, Ledebour 1842, Rogovich 1855, 1869, Chernyaev 1859, Herbich 1859, Knapp 1872, Montrezor 1886, 1887, 1888, 1890a, 1890b, Shmalhausen 1886, Paczoski 1897, 1899, 1900, 1911, Zapałowicz 1906, 1911a, 1911b). From the middle of the 19th century the authors began to indicate the origin of the alien species. The first such information is found in the works of P. Rogovich (1855) and I. Schmalhausen (1895-1897).

From the end of the 19th and beginning of the 20th century, due to the intensive farming of the steppes and the increase of arable lands, the focus was on the study of alien weeds. This direction had been actively developing in the 1920-1930s. Investigation of alien weeds, in addition to the usual morphological and agronomic characteristics, also included the analysis of the causes and ways of rapid resettlement of these plants, and peculiarities of their biology. The most informative among these publications are: the work by J. Paczoski (1911), on the distribution of *Amaranthus albus* L., by O. Yanata (1927) on the weeds of Ukraine, that includes the means of prevention, control and containment of their distribution, by A. Kuzmenko (1927) on

the distribution and biological features of *Fagopyrum tataricum* (L.) Gaertn., by O. Baranovskyi (1938) on the biology and control of *Galinsoga parviflora* Cav., etc. The general summary on the weeds of the country was published at 1937 under the editorship of E. Bordzilovsky. It contains information about a number of alien plants, their origin, and means of spreading in Ukraine.

V. Taliev (1900, 1902) was the first to note that anthropogenic activity had become a botanical and geographical factor. The anthropogenic factor, in his opinion, was determining the history of development and distribution of vegetation. His ideas stimulated the development of new perspectives on the introduction of alien plant species, as one of the aspects of modern changes in plant cover. In the 1920s a special study of alien plants as a special group began. M. Kotov investigated the mass resettlement of alien plants associated with human activity (1921, 1923, 1926, 1927a, 1927b–1929, 1934, 1949). In these works, he establishes the reasons for the appearance of alien plants in the country, the ways of their spreading, conditions that promote their distribution, etc. In parallel, the main floristic trend was developed. So, by the middle of the 20th century, large floristic material and datasets were collected, processed and published.

In the second half of the 20th century a comprehensive study of the regional alien flora of the Forest-Steppe, Steppe and Crimea was started with the works by V. Protopopova (1965) and S. Kozhevnikova (1970). In particular, their species composition was established, the structural analysis was conducted, and the degree of naturalization and regional differences were determined. In the study of the alien flora of forest-steppe and steppe zones (Protopopova 1965, 1973) particular attention is paid to the correlations between the distribution of alien plants and changes in the plant cover, determined by human activity. Based on numerous herbarium and literary data, the time of introduction and ways of resettlement of more than 100 species of alien plants from the end of the 18th century until the second half of the 20th century was established. The centers of introduction, ways of forming of the areals of these plants in the region, and analysis of the effectiveness of the means of distribution and biological characteristics are described. In particular, it is noted that anthropochory provides mainly the remoteness of new localities, and widespread depends on the effectiveness of natural methods of distribution of the propagule. Some taxa (*Xanthium* L., *Iva* L.) are revised, new species and forms are revealed. As a result of the historical reconstruction of the expansion of a number of species (*Galinsoga parviflora*, *Iva xanthiifolia* Nutt., *Amaranthus albus*, *A. blitoides* S. Watson, *Impatiens parviflora* DC., *Ambrosia artemisiifolia* L., *Solanum rostratum* Dunal., *Xanthium albinum* (Widder) H.Scholz, *Grindelia squarrosa* (Pursh) Dunal, *Cenchrus longispinus* (Hack.) Fernald, *Echinocystis lobata* (Michx.) Torr. & A. Gray, etc), it is shown that their distribution is conditioned by the influence of some favorable factors. In parallel, an inventory of the species of the alien fraction of Crimean flora was conducted. An annotated checklist of this group in the region's flora consisted of 133 species of vascular plants (Kozhevnikova, Rubtsov, 1971). In this work, its ways of introducing and distributing of the species of alien plants on the peninsula were considered in detail. The analysis of the geographical structure showed that the origin of the alien fraction of the Crimean flora is dominated by the species of the Ancient Mediterranean. Second place was occupied by North American species, most of which were weeds, and the third – East Asian species. The

bioecological structure of the fraction by the author based on the original material was characterized. By the degree of naturalization, the majority are ephemeroxytes, and the least number belong to neophytes. According to modern data, the alien fraction of the Crimean flora comprise 375 taxa; interesting information about the dynamics of its composition for almost 50 years and changes in the structure are given by N. Bagrikova (2013). In particular, only the numerosity of this group has increased by 242 species of vascular plants. According to the comprehensive analysis, the author established an increase in the modernization of flora, increased participation of Asian species, and in the spectrum of life forms – participation of trees (11.2%) and shrubs (6.9%). A specific feature of the alien fraction of the flora of Crimea is a high rate of the index of instability (0.56), which according to the author, is due to a significant number of ephemeroxytes (23.7%) and colonophytes (32.0%).

For the first time, the alien fraction of the flora of Ukraine was allocated as a part of synanthropic flora at the end of the 20th century; in total there were 646 species of vascular plants (Protopopova 1991) (Fig. 1). Structural analysis of the fraction was carried out and its characteristic features and regional peculiarities were determined. It was proven that the group of alien species generally caused changes to the systematic, geographical, and ecological structure of the flora of Ukraine and changes to the phytocenotic spectrum and spectrum of life forms, as well as expanded the florogenetic bonds with floras of the xerophytic regions of other continents. It was established that the distribution of alien plant species in Ukraine is characterized by certain patterns, which is evidenced by, for example, the calculated digital indicator of relative equilibrium of unstable and stable components in the species composition of the alien and synanthropic regional floras. A new version of the florogenetic classification of alien plants is proposed; in it the main criterion is the differentiation of species on the basis of their florogenetic bonds with some phytochorions within the climatic regions. This allowed us to establish that the main role in the formation of the alien fraction of the flora of Ukraine was played by the arid floras of the Ancient Mediterranean, North American prairie, Areas of Rocky Mountains, the Madrean Region of North America, and Chilean Patagonian arid regions of South America. That is, the process of adventization occurs through thermoxerophytization. Analysis of zonal-regional relationships in allocation of synanthropic species showed that everywhere anthropogenic localities are dominated in number by alien plants, especially in Crimea and the forest-steppe zone. Their number increases in the latitudinal direction to the south. However, their percentage in synanthropic flora increases in the opposite direction. Distribution of archeophytes and neophytes is characterized also by different trends: the number of archeophytes is increasing in the northern direction, and the neophytes – in the southern. Parameters of modernization and instability indexes also increase in the southern direction. It is shown that the unstable component of the alien fraction of flora (ephemeroxytes and ergasiophytes) do not affect the trends of zonal distribution of alien plants. The hypothesis of the origin of a number of narrow-endemic races of the regional flora from archeophytes in the process of their adaptation to new climatic conditions or ecotops is proposed.

One of the interesting trends in the investigation of the alien fraction of the flora of Ukraine of that time is the research of the anthropogenic transformation of ecosystems, first of all, their phytodiversity. For the first time, the reasoning of such a

research trend as an analysis of composition and structure of flora for a directed transformation of plant cover is proposed. Also, the method of flora-isolates (an elementary territorial unit of anthropogenic flora) is introduced. The first typologies of transformed flora and macroecotops of partial floras for the role in florogenesis have been developed. The functional role of various types of anthropogenic floras in reproduction and preservation of plant cover was substantiated. A significant role of phytinvasions in the processes of transformation of modern flora has been proved. These problems are best covered by an example of the structure, genesis, transformation and development of measures for the conservation of the flora of the southeast of Ukraine (Kondratyuk et al. 1980, 1985, 1988, Burda 1991).

In fact, the results that were published in the early 1990's, have intensified the development of the research of the regional alien flora at the end of the last century, which continues to this day. The alien fractions of the flora of some regions (Ostapko et al. 2009, 2010, Oitsius' 2011, Bagrikova 2013, Dvirna 2014, 2015, Kucher 2016, 2017, etc.), different floocomplexes (incl. railways and human-made ecotops (Burda & Tokhtar, 1992, Tokhtar 1993, 2005, Drel' 1997) and seaports (Petryk 1993), agroecosystems (Dzyuba 1990, 2000, Burda 2001, 2003, 2018, Mariyushkina 2003), urban floras (Mosyakin & Yavorska, 2002, Khlystun 2006, Zavyalova 2008, Baranovskiy et al. 2012), protected areas (Dubyna & Protopopova 1985, Protopopova et al. 1999, Shevchyk & Senchylo 2009, Melnychuk & Trokhymenko 2014, Burda et al. 2015, Zavialova 2017, etc.), and water reservoirs (Konogray 2008), etc., were investigated. During that time, there were numerous publications devoted to the finds of new species of aliene plants, which are collected in the bibliographic references (Burda et al. 2018). The amount of information about this group in general floristic publications has increased (Manual ..., 1987, Mosyakin & Fedoronchuk, 1999), in the checklists of the regional floras (Kondratiuk et al. 1985, Bairak 1997, Chopyk et al. 1998, Kucherevskiy 2003, Vasylyeva & Kovalenko, 2003, Tarasov 2005, 2012, Dzhuran et al. 2007, Kucherevskiy & Shol, 2009, Lukash 2008, Moysiienko 2011, etc.).

Among the recent regional studies, there are those in which the study of alien plants was one of the tasks. In particular, as a result of the study of plant contamination of the plant cover of the Middle Dnipro in its composition, 574 alien species were identified (Dzhuran et al. 2007). The adventization of the flora of Eastern Polissya is caused by 268 species of alien plants (Lukash 2008, 2009) and the Northern Black Sea Region – 565 (Moysiienko 2011). At the same time, there are works devoted exclusively to the comprehensive study of the alien species in the flora of some of the forest-steppe (Dvirna 2015) and steppe (Kucher 2017) territories of Ukraine. The specificity of both of the investigated floras is manifested in their mesophilic character, the general predominance of species with a wide range and wide ecological-coenotic amplitude, prevalence of a stable component over unstable, which in general corresponds to the characteristics of the alien fraction of the flora of Ukraine.

The similar comprehensive investigations of the alien fractions of regional floras allowed to find out the general features of the distribution of alien plants species, the peculiarities of the structure of this group and naturalization of species with the different origin, the causes of "outbreaks", the spreading of some species, etc.

Another important trend in the investigation of non-native plants in the flora of Ukraine is the special study of phytinvasions that was started at the beginning of the

21st century (Protopopova et al. 2002a, 2003). The composition of the fraction of 830 species of alien plants was established and the process of adventization of the flora over 150 years (1855-2002) was analyzed in these publications. In general, the progressive character of the adventization process, the tendency of increasing of the species number and the stepping up of the rates of introduction, distribution and naturalization, the outspread of ecotopological differentiation were generally noted. The participation of alien plants in the flora of Ukraine has increased from 3% (1855) to 15% (2018). Only 60% of species of the total fraction are successfully naturalized in all of the botanical and geographical zones of Ukraine. The role of species of alien plants in ecosystems according to the classification by J. Falinsky was evaluated. Most of the species have achieved biogeographic success, they have become widespread mainly in man-made areas with reduced competition. About 100 species of alien plants have become phytocoenotically successful: they have become established in semi-natural and natural plant communities. This has begun to affect their composition, structure, dynamics, and functional relationships. The least species of alien plants of the flora of Ukraine have acquired genetic success, that is, they are adapted mainly by hybridization (*Xanthium albinum*, *Centaurea diffusa* Lam., *Salix fragilis* L.).

The expansion of 30 species of alien plants, which took place during the second half of the 20th century, have been analyzed. It was found that in some species this process differs according to the continuance, the rate of distribution, or the character of development (outbreak, gradual stable distribution, fluctuations). Stress tolerance, a high degree of naturalization, effective means and a rapid rate of distribution, high coenotic activity, and wide ecological amplitude are the characteristics of all species in the state of expansion. There is a tendency of growth to the degree of naturalization and invasive ability for many species. As a result of the large amount of information on the changes caused by the widespreading and establishing of the species of alien plants in the plant communities, authors conclude that the adventization of the flora of Ukraine is increasing. Its impact is significant for the environment at the species, population, coenoses and ecosystem levels. Taking that into account, the working version of the National Strategy for the Control of Invasive Plant Species in Ukraine and supplementing the relevant national legislation has been proposed (Protopopova et al. 2002a, 2003).

At the same time, plant invasions in agroecosystems are investigated, according to the results of which, the threats of alien plants for agro-landscapes of Ukraine were evaluated (Burda 2001, 2003). Also, the ecological peculiarities in agroecosystems of Ukraine (Mariyushkina 2001, 2002a, 2002b, 2003), segetal potential (Burda 2003), and expansion of some species on the regional level were investigated (Lytvynenko & Vynokurov 2003–2004, Prots et al. 2011, Vykhov & Prots 2012, Kucher 2012, Dvirna 2014, etc.). The inventory of the species composition of alien plants in agricultural biotopes in Ukraine was conducted, in total 743 taxa were found (702 species, 24 subspecies and 17 hybrids). Some of them (129 species) are widespread in 49 countries of Europe. It has also been established that the species composition of alien plants, which is the biological contamination of agricultural biotopes in Ukraine, is similar to other European countries (Burda 2018).

In Ukraine, invasive plants were studied actively at different levels of organization, in particular, species, population, coenosis, biotope, etc. (Panchenko 2005,

Protopopova et al., 2009, Vykhor & Prots, 2012, 2014, Vykhor 2015, Didukh et al. 2016, Tokaryuk et al. 2018, etc.). The inventory of the species composition of alien plants at the different territorial levels is still relevant. At the beginning of this century, for the first time in Ukraine, an annotated list of species (94) of alien plants with a high invasive ability was proposed; the species displayed their biogeographic and phytocoenotic activity. Taking into account previous studies and the current state of phytodiversity in Ukraine, the species of vascular plants, most harmful for the phytodiversity on the Protected Areas of national importance, have been revealed. 100 invasive alien species occurring in Protected Areas of Ukraine are given as a brief compendium of three lists: Black (17), Grey (50) and Watch (33) (Zavyalova, 2017).

Some publications are devoted to the means of control of highly-invasive plant species (Maryushkina 1986, Mariyushkina & Hrytsenko 2000, Mariyushkina 2001, 2002a, 2002b, 2006, Petryk 2006, Borzykh 2014, etc.). As the results of investigation of 58 invasive species in the flora of the Northern Black Sea Region, their degree of naturalization, structural peculiarities, and participation in plant communities were established. The regional features of the group are a higher percentage of trees, shrubs and perennials, compared to the alien fraction of the total Ukrainian flora, as well as a prevalence of mesophytes, even though the region is situated in xerophilic conditions. These features can be explained by climate changes in recent decades, as a result of which it has become milder and more humid. The plant communities of some classes of native vegetation, such as, *Koelerio-Corynephoretea canescentis* Klika in Klika bet Novak 1941 (incl. *Festucetea vaginatae* Soó ex Vicherek 1972 (Mucina et al., 2016)), *Mollinio-Arrhenatheretea* Tx. 1937, *Festuco-Brometea* Br.-Bl. et Tx. ex Soó 1947, and others, have the highest invasibility. Invasive species have been noted in all types of habitats of the region, in particular *Centaurea diffusa* – in 6 anthropogenic and 4 native, *Conyza canadensis* (L.) Cronquist – in 7 and 3, *Grindelia squarrosa* – in 5 and 3 (Protopopova et al. 2009).

In the flora of Transcarpathia 43 species of invasive plants were found. Among them neophytes prevail, mostly those of North American origin; hemicryptophytes and therophytes, mesophytes, heliophytes (Prots 2015, Vykhor 2015). Dynamics trends of the distribution of invasive species in the flora of Transcarpathia and their ecological-cenototic confinement were analyzed. The distribution pattern of invasive species consists of a long lag-phase and an exponential phase. According to the study results, the growth of populations of invasive plants is slow in the lag-phase; the lower chronological limit of the exponential phase is established; the majority of the species are confined to the belt of lowland oak forests. Based on the study of spreading, the author developed predictive climate models of *Heracleum sosnowskyi* distribution. It is established, that the plants of the species can potentially populate all big riverside habitats, and the upper limit of their distribution will shift higher and reach the altitude of alpine meadows. This will be a threat to the structural and functional integrity of the high mountain river habitat types, meadows, subalpine and alpine areas. The assessment of the negative impact of invasive species, such as, *H. sosnowskyi*, *Helianthus tuberosus* L., *Reynoutria japonica* Houtt., *Solidago canadensis* L. was carried out (Vykhor 2015).

The invasive species of the flora of Bukovinian Cis-Carpathia region (29 species, including 5 transformers) was comprehensively studied. The current distribution was characterized, the cadastres and maps have been created, their participation in plant

communities has been analyzed, and a syntaxonomic scheme has been proposed (Tokaryuk et al. 2012, 2018).

The nonnative species as well as transformers were studied in different regions of Ukraine: Transcarpathia, Bukovinian Cis-Carpathia, the Middle Dnipro Region, Ukrainian Polissya, the Northern Black Sea Region, the South Coast of Crimea, Romensko-Poltavsky geobotanical district and Starobilsk grass-meadow Steppe (Protopopova et al. 2009, 2010a, 2012, 2014a, 2015, Dvirna 2014, Kucher 2015). According to the results of these investigations the different composition of this group has been detected. The common features of the transformers of all studied areas were established. Among them, the prevalence of alien species with their origin from America or Eastern Asia, mesophytes – in ecological spectra, the higher percentages of trees or shrubs were revealed. The prevalence of species with Mediterranean origin and plants from the xerothermic habitats in the composition of transformers of the South Coast of Crimea has been noted.

The ergasiophytes of the flora of Ukraine (458 species, which comprised about half of its alien fraction (46.5%) and every time a new ones appear) were analyzed (Protopopova & Shevera 2014). There is the prevalence of the species of Mediterranean and North American origin in the composition of this group. A high percentage of trees and shrubs in the spectrum of life forms and mesophytes in the hygromorphs spectrum of ergasiophytes of the flora of Ukraine were revealed. By the degree of naturalization [according to the classification by A. Thellung (1915)], 78.8% of species are ergasiofigophytes. There was an increase in the degree of naturalization in 23.2% of species over the last 150 years. There are transformers in this group. The spreading of some ergasiophytes as a result of population growth has reached expansion (there are 20-120 years from the first records to expansion). High levels of propagule pressure some of ergasiophytes as a result of planting lead to species eventually escaping from cultivation, regardless of biological traits.

One of the trends of the study of alien plants of the flora of Ukraine is a special comprehensive research of the urban floras in the different botanical and geographical zones of Ukraine initiated by R.I. Burda (1982) and has been conducted since the late 1980's. The urban floras of Kherson (Moysienko 1999), Mykolaiv (Melnyk 2001), Uzhhorod (Protopopova & Shevera 2003), Kamyianets-Podilskiy (Kagalo et al. 2004), Kryvyi Rig (Shol 2004), Netishyn, Ostrog, Slavuta, Shepetivka (Gubar 2006), Kirovograd (now Kropyvnytskyi) (Arkushina 2007), Simferopol (Epikhin 2008), Lutsk, Chernihiv (Zavyalova 2012), Kuznetsovsk, Dubrovytsia, Sarny, Kostopil, Berezne (Hutsman 2013), and urban area of the Donetsk-Makiivka (Derevenska 2014), Kharkiv (Zviagintseva 2015), etc. The results of this investigation have found the portion of the alien fraction of the studied urban floras in Ukraine is approximately the same and does not exceed 28% of the total species composition (Fig. 2).




Fig. 2. The total species richness of the urban floras of Ukraine and shares of their alien fractions

The annotated checklists of the urban floras of Donetsk, Luhansk, Mariupol, Sloviansk (Burda 1997), Uzhhorod (Protopopova & Shevera 2002), Kryvyi Rig (Kucherevskiy & Shol 2003, 2009), Kamianets-Podilskiy (Kagalo et al. 2004), Lutsk, Kirovograd (Arkushina & Popova 2010), Pereyaslav-Khmelnytskyi (Fedoronchuk et al. 2010), Chernihiv (Zavyalova 2010), Kharkiv (Zviagintseva 2015) or some of these fractions, such as, the synanthropic flora of Odesa (Vasylyeva-Nemertsalova 1996), the nonnative flora of the Kyiv urban area (Mosyakin & Yavorska 2002), etc., were published. As the results of these studies of the urban floras of Ukraine the general and regional features were revealed. Such as, highest species richness, the high levels of the synanthropization (46.7%–68.6%) and adventization, extension of habitats differentiation, the changes of structure in comparison with regional floras, the leveling of zonal peculiarities, for example, the mesophytization of the floras of Steppe zone cities, S and V-similar character of distribution of the species, etc. (Vasylyeva-Nemertsalova 1996, Moysyenko 1999, Shevera 2005, Zavyalova 2012). The comparative analysis of the species composition of neophytes and archeophytes of 12 urban floras using the mathematical statistics methods are provided. The peculiarities of their distribution, in particular, of zonal-regional conditions, and character of anthropogenic impact, and socio-economic or historical factors have been detected (Protopopova & Shevera 2008b, Protopopova et al. 2010, 2013). Methods of study of

adaptive strategy of alien plants on urban territory was generalized by R. Burda & O. Ignatyuk (2011).

The adventization of the plant cover of the Protected Areas of Ukraine began to be studied in the 1980s. One of the first studies in this trend was the investigation of the alien fraction of the floras of the “Mys Martyan” Nature Reserve (Golubyeva 1982) and Danube Nature Reserve (Dubyna & Protopopova 1985). In fact, the total species composition of the fraction, its structure and regional peculiarities have been investigated. After that, similar studies of the alien species in the floras were carried out in the National Nature Parks “Synevyr” (Protopopova et al. 1999), “Vyzhnytsky” (Chorney et al. 2005), Kaniv Nature Reserve (Shevchyk & Senchylo 2009), protected areas of the Kerch Peninsula in the Crimea (Bagrikova 2011), National Nature Park “Biloberezhzha Sviatoslava” (Melnychuk & Trokhymenko 2014), protected areas of the Forest-Steppe of Ukraine (Burda et al. 2015), etc. Some data about alien plant species in the floras of the Protected Areas consist of its checklists (Bioriznomanittia Karpatskoho ..., 1997, Bioriznomanitnist Dunayskoho..., 1999, Nechytaylo et al. 2002, Oliyar 2002, Dubyna et al. 2003, Panchenko 2005, Derzhypilskiy et al. 2006, Kostina & Bagrikova 2010, Kolomiychuk & Yaroviyi 2010, Orlov 2013, Karpenko, 2016, Baranovskiy et al. 2017, etc.). The biological contamination of the National Econet began to study in the 2013th (Zavyalova & Korniyenko 2014).

For the first time the state of the biological contamination of the floras of Protected Areas of the Forest Steppe of Ukraine have been summarized early in the 21st century. The investigation of alien invasive species with an assessment of their environmental impact were carried out for 14 local floras of the forest-steppe zone (Burda et al. 2015). A total species composition (354 taxa: 345 species, 2 subspecies and 7 hybrids) and 17 transformers was established. According to the authors' data the peculiarities of the life forms spectrum is the prevalence of therophytes and a high proportion of phanerophytes. The features of the non ability of ecosystems of the Protected Areas for resistance to the threats of phyto-invasions have been defined. According to the assessment of the impact of alien plants to the plant cover of the investigated objects (with the exception of *Salix fragilis*) its reverse character was revealed (Burda et al. 2015). At the same time, the study of invasive alien species in the plant cover of Protected Areas of national importance to Ukraine was conducted and they are given as a brief compendium of three lists: Black, Grey and Watch. These lists were compiled on the basis of set criteria. The participation of the studied species in the composition of plant communities was established as well as their representation in the territory of more than 50 protected areas of Ukraine of different categories. We have found that about a third of the studied species have had a significant impact on the species, coenotic and ecosystem levels, and threatening the native biodiversity of the Protected Areas of Ukraine. The rest is a potential threat. At the same time, some alien species of the Grey and Watch lists threaten native biodiversity at local or regional level in some native areas of Ukraine. They are dangerous only for certain types of natural habitats or plant communities of certain vegetation types (Zavialova 2017).

The main results of the critical taxonomic revisions, including of alien plant species, are available in multi-volume publications, manuals of plants, monographs and articles: Flora of the RSS Ucr. (1935-1965), Flora URSS (1934-1964), Manual of vascular plants of Ukraine (1950, 1965, 1987), Buriyany Ukrainy (1970), Flora Partis

Europaeae URSS (1974-1996), Flora Europaeae Orientalis (1996–2004), Ecoflora of Ukraine (Didukh et al. 2000-2010), Zlaki Ukrainy [Grasses of the Ukraine] (Prokudin et al. 1977), The genus *Oenothera* L. in Eastern Europe (Rostański et al. 2004), and many other taxa (Protopopova 1964, 1974, Mosyakin 1988, 1989, 1992, 1995a, b, Mosyakin & Korniyenko 2006, etc.). Mostly, the critical taxonomic study was concerned with the alien plant species of the families of Asteraceae Dumort., Poaceae Barnhart, Brassicaceae Burnett, Cucurbitaceae Juss., Malvaceae Juss., Boraginaceae Juss., Solanaceae Juss., Amaranthaceae Juss., Balsaminaceae A. Rich., etc.

Conclusion

During the analyzed period of study of the alien fraction in the flora of Ukraine, rich and diverse factual material has been accumulated and summarized. Thereby, a large many-sided or various factual material on different trends of research for the analyzed period of the study of the alien fraction of the flora of Ukraine. Its species composition at different territorial levels: national, regional and local, which significantly differs quantitatively and qualitatively as a result of the constant gain of the adventization process have been established. The checklists of the alien fraction of the flora of Ukraine, and numerous regional and urban floras etc., have been consisted. A group of invasive species, including transformers, have been dedicated and researched. Some characteristics of non-native species in the plant cover of the Protected Areas of national importance of Ukraine, in particular, the composition, representation, participation, impact etc., have been investigated. The populations of some alien species have been researched. Some practical aspects of the problem, such as, bioecological features, segetal potential of weeds and its role in agroecosystems have been included to the investigations too. Consequently, these and other studies represent a great variety of floristic, chorological, ecological, phytocoenotical, taxonomical, and other directions of comprehensive scientific work in Ukraine, the results of range from the first records of non-native species of plants to the theoretical generalizations and conclusions about the peculiarities of the alien fraction of the flora formation in the country.

References

- Andrzejowski, A. (1823). Rys botaniczny krain zwiedzonych w podrózach pomiędzy Bohem a Dniestrem od Zbrucza aż do Morza Czarnego odbytych w latach 1814, 1816, 1818, 1822. Wilno, V(1): 1–126. (In Polish).
- Andrzejowski, A. (1860). Ischislenie rasteniy Podolskoy gubernii i smezhnykh s neyu mest. I. *Tr. Komis. pri un-te sv. Vladimira dlya opisaniya guberniy Kievsk. uchebn. okruga*, V(1): 1–51. [Анжейовский, А.Л., 1860 – Исчисление растений Подольской губернии и смежных с нею мест. I. *Тр. Комис. при ун-те св. Владимира для описания губерний Киевск. учебн. округа*, V(1): 1–51]. (In Russian).
- Andrzejowski, A. (1862). Ischislenie rasteniy Podolskoy gubernii i smezhnykh s neyu mest. II. *Tr. Komis. pri un-te sv. Vladimira dlya opisaniya guberniy Kievsk. uchebn. okruga*, 7: 94–142. [Анжейовский, А.Л., 1862 – Исчисление растений Подольской губернии и смежных с нею мест. II. *Тр. Комис. при*

- ун-те св. Владимира для описания губерний Киевск. учебн. округа, 7: 94–142]. (In Russian).
- Arkushina, H.F. (2007). Urbanoflora Kirovohrada: avtoref. dys. ... kand. biol. nauk. 20 s. Yalta. [Аркушина, Г.Ф., 2007 – Урбанофлора Кіровограда: автореф. дис. ... канд. біол. наук. 20 с. Ялта]. (In Ukrainian).
- Arkushina, H.F., Popova, O.M. (2010). Konspekt flory sudynnykh roslyn m. Kirovohrada. 232 s. Kirovograd: Polimed-Servis. [Аркушина, Г.Ф., Попова, О.М., 2010. Конспект флоры судинных растений м. Кіровограда. 232 с. Кіровоград: Полімед-Сервіс]. (In Ukrainian).
- Bagrikova, N.A. (2011). Analiz adventivnoy fraktsii flory prirodnykh zapovednikov Kerchenskogo poluostrova (Krym). *Ekosistemy, ikh optimizatsiya i okhrana*, 4 (23): 27–34. [Багрикова, Н.А., 2011 – Анализ адвентивной фракции флоры природных заповедников Керченского полуострова (Крым). *Экосистемы, их оптимизация и охрана*, 4 (23): 27–34]. (In Russian).
- Bagrikova, N.A. (2013). Strukturnyi analiz adventivnoy fraktsii flory Krymskogo poluostrova (Ukraina). *Ukr. Bot. J.*, 70(4): 489–507 [Багрикова, Н.А., 2013 – Структурный анализ адвентивной фракции флоры Крымского полуострова (Украина), *Укр. ботан. журн.*, 70(4): 489–507]. (In Russian).
- Baranovskiy, V.O., Maniuk, V.M., Ivan'ko, I.A., Karmyzova, L.O. (2017). Analiz flory Natsionalnoho pryrodnoho parku "Orilskiy", 320 s. Dnipro: Lira press. [Барановський, В.О., Манюк, В.М., Іванько, І.А., Кармизова, Л.О., 2017 – Аналіз флори Національного природного парку "Орільський", 320 с. Дніпро: Ліра]. (In Ukrainian).
- Baranovskiy, V.O., Tarasov, V.V., Ivan'ko, I.A., Karmyzova, L.O. (2012). Adventyvena flora m. Dnipropetrovsk. In: Synantropizatsiya roslynnoho pokryvu Ukrainy: tezy naukovykh dopovidey (27–28 veresnia 2012 r.), p. 14–17. Kyiv & Pereyaslav-Khmelnyskiy. [Барановський, В.О., Тарасов, В.В., Іванько, І.А., Кармизова, Л.О., 2012 – Адвентивна флора м. Дніпропетровська // Синантропізація рослинного покриву України: Тези наук. доп. (27–28 вересня 2012 р.), с. 14–17, Київ, Переяслав-Хмельницький.]. (In Ukrainian).
- Baranovskiy, O.L. (1938). Biolohiya bur'yanu *Galinsoga parviflora* Cav. ta zasoby borotby z nym. *Pratsi Zhytomyrskoho silsko-hospodarskoho instytutu*, 1: 3–94. [Барановський, О.Л., 1938 – Біологія бур'яну *Galinsoga parviflora* Cav. та засоби боротьби з ним. *Пр. Житомир, с.-г. ін.-ту*. 1: 3–94]. (In Ukrainian)
- Bayrak, O.M. (1997). Konspekt flory Livoberezhnoho Prydniprov'ya. Sudynni roslyny / Red. S.L. Mosyakin. 164 s. Poltava: Verstka. [Байрак, О.М., 1997 – Конспект флоры Лівобережного Придніпров'я. Судинні рослини / Ред. С.Л. Мосякін. 164 с. Полтава: Верстка]. (In Ukrainian).
- Besser, W. (1809). *Primitiae florum Galiciae austriacae utriusque*. Vol. 1–2, 822 p. Vienna. (In Latin).
- Besser, W. (1822). *Enumeratio plantarum hucusque in Volhynia, Podolia, Gub. Kiioviensi, Bessarabia cisthyraica et circa Odessam collectarum, simul cum observationibus in Primitias Florae Galiciae Austriacae*. 111 p. Vilnae. (In Latin).
- Bieberstein, M.F. (1808a, b, c). *Flora Taurico-Caucasica*. Vol. 1, 428 p., vol. 2, 478 p., vol. 3, 336 p. Charkovia. (In Latin).

- Bioriznomanitnist Dunayskoho biosferneho zapovidnyka, zberezheniya ta upravlinnya (1999). Red. Yu.R. Shelyah–Sosonko. 703 s. Kyiv: Interekotsentr. [Біорізнманітність Дунайського біосферного заповідника, збереження та управління, 1999 – Ред. Ю.Р. Шеляг–Сосонко. 703 с. Київ: Інтерекоцентр]. (In Ukrainian).
- Bioriznomanittia Karpatskoho biosferneho zapovidnyka (1997). Red. rada Ya.I., Movchan, F.D., Hamor, Yu.R., Shelyag-Sosonko, I.O., Dudka, I.V., Zagorodniuk. 715 s. Kyiv: Interekocenter. [Біорізнманіття Карпатського заповідника, 1997 – Ред. рада Я.І., Мовчан, Ф.Д., Гамор, Ю.Р., Шеляг–Сосонко, І.О., Дудка, І.В., Загороднюк. 715 с. Київ: Інтерекоцентр]. (In Ukrainian).
- Borzykh, O.I. (2014). Kontrol poshyrenykh vydiv bur'yaniv iz vohnyshchamy hirchaka povzuchoho *Acroptilon repens* L. Karantyn i zakhyst Roslyn, 1: 17–18 [Борзих, О.І., 2014 – Контроль поширених видів бур'янів із вогнищами гірчака повзучого *Acroptilon repens* L. Карантин і захист росл. 1: 17–18]. (In Ukrainian).
- Visyulinoi, O.D., Kotova, M.I., Dobrochayevoi, D.M. & Kuznyetsovoi, H.O. (red.) (1970). Buriyany Ukrainy. Kyiv: Naukova dumka. [Бур'яни України, 1970 – Під заг. ред. О.Д., Вісюліної, М.І., Котова, Д.М., Доброчасвої, Г.О., Кузнецової. Київ: Наукова думка]. (In Ukrainian).
- Burda, R.I. (1982). Urbanoflora kompleksa Donetsk–Makeevka. Pp. 11–12. In VII syezd Ukr. botan. o-va: Tez. dokl. Kiev: Naukova dumka. [Бурда, Р.І. (1982). Урбанофлора комплексу Донецьк–Макеевка. В кн. VII съезд Укр. ботан. о-ва: Тез. докл. С. 11–12. Киев: Наук. думка]. (Un Russian).
- Burda, R.I. & Tokhtar V.K. (1992). *Cenchrus pauciflorus* Benth. na Pivdennomu Skhodi Ukrainy. *Ukr. botan. zhurn.* 49 (5): 14–18. [Бурда, Р.І., Тохтар, В.К. (1992). *Cenchrus pauciflorus* Benth. на Південному Сході України. *Укр. ботан. журн.* 49(5): 14–18]. (In Ukrainian).
- Burda, R.I. (1991). *Antropogennaya transformatsiya flory*. 168 s. Kiev: Naukova dumka [Бурда, Р.І., 1991 – Антропогенная трансформация флоры. 168 с. – Киев: Наук. думка]. (In Russian)
- Burda, R.I. (1997). *The checklist of the Donbass urban flora*. 50 p. Donetsk.
- Burda, R.I. (2001). Otsinka ekolohichnoi zahrozy zanosnykh roslyn v ahrolandshaftakh Ukrainy. *Promysh. botan.: Sbornik nauch. trudov.* 1: 16–21 [Бурда, Р.І., 2001 – Оцінка екологічної загрози заносних рослин в агроландшафтах України. *Промыш. ботан.:* Сб. науч. тр. 1: 16–21]. (In Ukrainian).
- Burda, R.I. (2003). Segetalniy potentsial zanosnykh rasteniy v agrolandshaftakh Ukrainy In: Problemy izucheniya adventivnoy i sinantropnoy flory v regionakh SNG. Mat. nauch. konf. / Red. V.S. Novikov i A.V. Shcherbakov. S. 29–30. Moskva: Izd-vo Botan. sada MGU; Tula: Grif i K0. [Бурда, Р.І., 2003 – Сеgetальний потенціал заносних рослин в агроландшафтах України. В кн.: Проблемы изучения адвентивной и синантропной флоры в регионах СНГ: Мат. науч. конф. / Ред. В.С. Новиков и А.В. Щербаков. С. 29–31. М.: Изд-во Ботан. сада МГУ; Тула: Гриф и К0]. (In Russian).
- Burda, R.I. (2018). Alien Plant Species in the agricultural habitats of Ukraine: diversity and risk assessment. *Ekológia*, 37(1): 24–31.
- Burda, R.I., Ignatyuk, O.A. (2011). Metodyka doslidzhennya adaptivnoi stratehii chuzhoridnykh vydiv roslyn v urbanizovanomu seredovyschi. 112 s. Kyiv: ZAT

- “Vipol” [Бурда, Р.І., Ігнатюк, О.А., 2011 – Методика дослідження адаптивної стратегії чужорідних видів рослин в урбанізованому середовищі. 112 с. Київ: ЗАТ “Віпол”] (In Ukrainian).
- Burda, R.I., Pashkevych, N.A., Boyko, H.V., Fitsaylo, T.V. (2015). Chuzhoridni vydy okhoronnykh flor Lisostepu Ukrainy. 118 s. Kyiv: Naukova dumka [Бурда, Р.І., Пашкевич, Н.А., Бойко, Г.В., Фіцайло, Т.В., 2015 – Чужорідні види охоронних флор Лісостепу України. 118 с. Київ: Наук. думка]. (In Ukrainian).
- Burda, R.I., Protopopova, V.V., Shevera, M.V., Kucher, O.O. (2018). *Chuzhoridni vydy flory Ukrainy: roky i avtory. Bibliohrafichnyi pokazhchuk*. Vypusk 5. 130 s. Kyiv [Бурда, Р.І., Протопопова, В.В., Шевера, М.В., Кучер, О.О., 2018 – Чужорідні види флори України: роки і автори. Бібліографічний покажчик. Вип. 5. 130 с. Київ]. (In Ukrainian).
- Chernyayev, V.M. (1859). *Konspekt rasteniy, dikorastushchikh i razvodimykh v okrestnostyakh Kharkova i v Ukraine*. 90 s. Kharkov [Черняев В.М., 1859 – Конспект растений, дикорастущих и разводимых в окрестностях Харькова и в Украине, 90 с. Харьков]. (In Russian).
- Чорук, В.І., Бортняк, М.М., Воутюк, Ю.О., Погребенник, В.П., Кучерява, Л.Ф., Нечитайло, В.А., Любченко, В.М., Шевчук В.Л. (1998). *Konspekt flory Serednoho Prydniprov'ya. Sudynni roslyny*. 140 s. Kyiv: Fitosotsiotsentr. [Чопик, В.І., Бортняк, М.М., Войтюк, Ю.О., Погребенник, В.П., Кучерява, Л.Ф., Нечитайло, В.А., Любченко, В.М., Шевчик В.Л., 1998 – Конспект флори Середнього Придніпров'я. Судинні рослини. 140 с. Київ: Фітосоціоцентр]. (In Ukrainian).
- Chorney, I.I., Budzhak, V.V., Yakushenko, D.M., Korzhyk, V.P., Solomakha, V.A., Sorokan, Yu.I., Tokaryuk, A.I., Solomakha, T.D. (2005). *Natsionalnyi pryrodnyi park "Vyzhnytskyi". Roslynniy svit*. Vyp. 4. 129–133. Kyiv: Fitosotsiotsentr. [Чорней, І.І., Буджак, В.В., Якушенко, Д.М., Коржик, В.П., Соломаха, В.А., Сорокан, Ю.І., Токарюк, А.І., Соломаха, Т.Д. 2005 – Національний природний парк “Вижницький”. Рослинний світ. Вип. 4. 129–133. Київ: Фітосоціоцентр.]. (In Ukrainian).
- Derevenska, H.H. (2014). *Urbanoflora ahlomeratsii Donetsk-Makiivka: avtoref. dys. ... kand. biol. nauk*. 24 s. Yalta. [Дерев'янська, Г.Г., 2014 – Урбанофлора агломерації Донецьк-Макіївка: автореф. дис. ... канд. біол. наук. 24 с. Ялта.]. (In Ukrainian).
- Derzhypilskyi, L.M., Tomych, M.V., Yusup, S.V. et al. (2006). *Natsionalnyi pryrodnyi park "Gutsulshchyna"*. Roslynniy svit / Eds. V.A. Solomakha, I.I. Chorney. Vyp. 9. 360 s. Kyiv: Fitosociocenter. [Держипільський, Л.М., Томич, М.В., Юсип, С.В. та ін., 2006 – Національний природний парк «Гуцульщина». Рослинний світ. Вип. 9 / Ред. В.А. Соломаха, І.І. Чорней. 360 с.– Київ: Фітосоціоцентр]. (In Ukrainian).
- Didukh, Ya.P. (red.) (2000–2010). *Ecoflora of Ukraine*. Vol. 1–3, 5–6, Kyiv: Fitosociocenter [Екофлора України / Відпов. ред. Я.П., Дідух. 2000–2010 – Т. 1–3, 5–6, Київ: Фітосоціоцентр]. (In Ukrainian).
- Didukh, Ya.P., Chorney, I.I., Budzhak, V.V., Tokaryuk, A.I., Kish, R.Ya., Protopopova, V.V., Shevera, M.V., Kozak, O.M., Kontar, I.S., Rozenblit, Yu.V., Norenko, K.M. (2016). *Klimatohenni zminy roslynnoho svitu Ukrainskykh Karpat / Red.*

- Ya.P. Didukh, I.I. Chorney. 284 s. Chernivtsi: Druk Art. [Дідух, Я.П., Чорней, І.І., Буджак, В.В., Токарюк, А.І., Кіш, Р.Я., Протопопова, В.В., Шевера, М.В., Козак, О.М., Контар, І.С., Розенбліт, Ю.В., Норенко, К.М., 2016 – Кліматогенні зміни рослинного світу Українських Карпат / Ред. Я.П. Дідух, І.І. Чорней. 284 с. Чернівці: Друк Арт.]. (In Ukrainian).
- Drel', V.F. (1997). *Adventywna flora zaliznyts Luhanskoyi oblasti: avtoref. dys. ... kand. biol. nauk.* 20 s. Kyiv. [Дрель, В.Ф., 1997 – Адвентивна флора залізниць Луганської області: автореф. дис. ... канд. біол. наук. 20 с. Київ.]. (In Ukrainian).
- Dubyna, D.V., Protopopova, V.V. (1985). Analiz adventyvnoy flory zapovednika "Dunayskiye plavni". *Biol. nauki*, 10: 68–73. [Дубына, Д.В., Протопопова, В.В., 1985 – Анализ адвентивной флоры заповедника Дунайские плавни, *Биол. науки*, № 10: 68–73]. (In Russian).
- Dvirna, T.S. (2014). Adventivnaya frakciya flory Romensko-Poltavskogo geobotanicheskogo okruga: analiz i konspekt. *Fitoraznoobraziye Vostochnoy Evropy*, 8(1): 4–19. [Двирна, Т.С., 2014 – Адвентивная фракция флоры Роменско-Полтавского геоботанического округа: анализ и конспект. *Фиторазнообразие Вост. Евр.*, 8(1): 4–19]. (In Russian).
- Dvirna, T.S. (2015). *Adventywna fraktsiya flory Romensko-Poltavskoho heobotanichnoho okruhu ta yiyi invazyynni potentsial: avtoref. dys. ... kand. biol. nauk.* 18 s. Kyiv. [Двирна, Т.С., 2015 – Адвентивна фракція флори Роменсько-Полтавського геоботанічного округу та її інвазійний потенціал: автореф. дис. ... канд. біол. наук. 18 с. Київ.]. (In Ukrainian).
- Dzhuran, V.M., Kretsul, N.I., Protopopova, V.V., Fedoronchuk, M.M., Shevera, M.V. (2007). *Fitozabrudnennya roslynnoho pokryvu Serednoho Prydniprov'ya. Antonovanyi konspekt synantropnoi flory.* 48 s. Kyiv: Avtoreferat [Джуран, В.М., Крецул, Н.І., Протопопова В.В., Федорончук, М.М., Шевера, М.В., 2007 – Фітозабруднення рослинного покриву Середнього Придніпров'я. Антонований конспект синантропної флори. 48 с. Київ: Автореферат.]. (In Ukrainian).
- Dzyuba, T.P. (1990). *Flora i rastitel'nost risovykh system Prichernomor'ya: avtoref. dis. ... kand. biol. nauk.* 20 s. Kiev. [Дзюба, Т.П., 1990 – Флора и растительность рисовых систем Причерноморья: автореф. дис. ... канд. биол. наук. 20 с. Киев.]. (In Russian).
- Dzyuba, T.P. (2000). Adventywna flora rysovykh system Prychornomor'ya: istoriya doslidzhennya ta osnovni problemy. In.: *Fitosotsiologiya. 100 rokiv naukovoho napryamku.* Mat. nauk. konf. (Kyiv, 1–3 zhovtnya 2000). S. 15-17. Kyiv: Fitosotsiotsentr. [Дзюба, Т.П., 2000 – Адвентивна флора рисових систем Причорномор'я: історія дослідження та основні проблеми. В кн.: Фітосоціологія. 100 років наукового напрямку. Мат. наук. конф. (Київ, 1–3 жовтня 2000). С. 15-17. Київ: Фітосоціоцентр]. (In Ukrainian).
- Epihin, D.V. (2008). *Sovremennoe sostoyanie rastitelnogo pokrova goroda Simferopol'ya: avtoref. dis. ... kand. biol. nauk.* 20 s. Yalta. [Епихин, Д.В., 2008 – Современное состояние растительного покрова города Симферополя: автореф. дис. ... канд. биол. наук. 20 с. Ялта.]. (In Russian).
- Fedoronchuk, M.M., Protopopova, V.V., Shevera, M.V., Dzhuran, V.M., Kretsul, N.I., Chorna, H.A., Hubar, L.M. (2010). Sudynni roslyny. In: *Pereyaslav-Khmelnyskiy. Pryroda: roslynniyi svit. Krytychniyi inventaryzatsiynyi anotovanyi*

- konspekt flory ta roslynnosti: sudynni roslyny, mokhopodibni, lyshaynyky, vodorosti*. S. 33–111. Korsun-Shevchenkivskiy: FOP Maydachenko I.S. [Федорончук, М.М., Протопопова, В.В., Шевера, М.В., Джуран, В.М., Крецул, Н.І., Чорна, Г.А., Губарь, Л.М., 2010 – Судинні рослини. В кн.: Переяслав-Хмельницький. Природа: рослинний світ. Критичний інвентаризаційний анований конспект флори та рослинності: судинні рослини, мохоподібні, лишайники, водорості / С. 33–111. Корсунь-Шевченківський: ФОП Майдаченко І.С.]. (In Ukrainian).
- Flora Europaeae Orientalis, 1996–2004 – Vol. 9–11, Moskva – Sankt-Peterburg: Mir i Semia-95; Tovaryshestvo nauchnykh izdaniy KMK [Флора Восточной Европы, 1996–2004 – Т. 9–11, Москва–Санкт-Петербург: Мир и семья-95; Товарищество научных изданий КМК]. (In Russian).
- Flora Partis Europaeae URSS, 1974–1994 – Vol. 1–8. Leningrad; Sankt-Peterburg: Nauka. [Флора Европейской части СССР, 1974–1994 – Т. 1–8, Ленинград; Санкт-Петербург: Наука]. (In Russian).
- Flora URSS Ucr., 1936–1965 – Vol. 1–12. Kyiv: Vydavnytstvo AN URSS. [Флора УРСР в 12-ти тт., 1936–1965 – Т. 1–12. Київ: Видавництво АН УРСР]. (In Ukrainian).
- Flora URSS, 1934–1964 – Vol. 1–30. Leningrad: Izdatelstvo AN USSR [Флора СССР в 30-ти тт., 1934–1964 – Т. 1–30. Ленинград: Издательство АН СССР]. (In Russian).
- Golubeva, I.V. (1982). Ob adventivnykh rastenyakh zapovednika “Mys Martyan”, *Bull. Nikitskyi Botan. Sada*, 49: 13–16 [Голубева, И.В., 1982. Об адвентивных растениях заповедника «Мыс Мартьян», *Бюл. Никит. ботан. сада*, 49, 13–16]. (In Russian).
- Gubar, L.M. (2006). *Urbanoflory skhidnoyi chastyny Maloho Polissia: avtoref. dys. ... kand. biol. nauk*. 20 s. Kyiv. [Губарь, Л.М., 2006 – Урбанофлора східної частини Малого Полісся: автореф. дис. ... канд. біол. наук. 20 с. Київ.]. (In Ukrainian).
- Güldenstädt, J.A. (1787). *Reisen durch Russland und im Caucasischen Gebürge. Auf Befehl der Russisch-Kayserlichen Akad. der Wiss. hrsg. von P. S. Pallas*. Theil 1, 511 S., 13 Einlag: Fig. St. Petersburg. (In German).
- Güldenstädt, J.A. (1791). *Reisen durch Russland und im Caucasischen Gebürge. Auf Befehl der Russisch-Kayserlichen Akad. der Wiss. hrsg. von P. S. Pallas*. Theil 2, 552 S., 14 Einlag: Fig. St. Petersburg. (In German).
- Herbich, F. (1859). *Flora der Bukowina*. 460 s. Leipzig. (In German).
- Hutsman, S.V. (2013). *Flora mist skhidnoi chastyny Volynskoho Polissya: avtoref. dys. ... kand. biol. nauk*, 20 s. Kyiv. [Гуцман, С.В., 2013 – Флора міст східної частини Волинського Полісся: автореф. дис. ... канд. біол. наук. 20 с. Київ.]. (In Ukrainian).
- Kagalo, O.O., Skibitska, N.V., Lyubinska, L.H., Guzik, J., Protopopova, V.V., Shevera, M.V. (2004). Sudynni roslyny m. Kamyantsia-Podilskoho. In: *Bioroznomanittia Kamyantsia-Podilskoho. Poperedniy krytychniy inventaryzatsiyniy konspekt roslyn, hrybiv i tvaryn*. P. 82–134. Lviv: Liga-Press [Кагало, О.О., Скібіцька, Н.В., Любінська, Л.Г., Гузік, Я., Протопопова, В.В., Шевера, М.В., 2004 – Судинні рослини м. Кам'янець-Подільського. В кн.: Біорізноманіття Кам'янець-Подільського. Попередній критичний інвентаризаційний конспект рослин, грибів і тварин. С. 82–134. Львів: Ліга-Прес, 2004]. (In Ukrainian).

- Karpenko, Yu.O. (2016). *Sudyinni roslyny Mezynskoho natsionalnoho pryrodnoho parku*. 100 s. Chernihiv: Desna Polihraf. [Карпенко, Ю.О., 2016 – Судинні рослини Мезинського національного природного парку. 100 с. Чернігів: Десна Поліграф.]. (In Ukrainian).
- Khlystun, N.Ya. (2006). *Adventywna flora m. Chernivtsi: avtoref. dys. ... kand. biol. nauk*. 20 s. Kyiv [Хлистун Н.Я., 2006 – Адвентивна флора м. Чернівці: автореф. дис. ... канд. біол. наук. 20 с. Київ]. (In Ukrainian).
- Knapp, J.-A. (1872). *Die bischer bekannten Pflanzen Galiciens und der Bucowina*. 267 s., Wien. (In German).
- Kolomiychuk, V.P., Yarovy, S.O. (2010). *Konspekt flory Pryazovskoho NPP*. 296 s. Kyiv: Alterpres [Коломійчук, В.П., Яровий С.О., 2011 – Конспект флори Приазовського НПП. Київ: Альтерпрес]. (In Ukrainian).
- Kondratyuk, E.N., Burda, R.I., Chupryna, T.G., Khomyakova, T.T. (1988). *Luganskyi gosudarstvennyi zapovednik. Rastitelnyi mir* / Ed. V.P. Tarabrin, 188 s. Kiev: Naukova dumka press. [Кондратюк, Е.Н., Бурда, Р.И., Чуприна, Т.Т., Хомякова М.Т. Луганский государственный заповедник. Растительный мир / Ред. В.П. Тарабрин. 188 с. Киев: Наук. думка. (In Russian)
- Kondratyuk, E.N., Burda, R.I., Ostapko, V.M. (1985). *Konspekt flory Yugo-Vostoka Ukrainy*. 272 s. Kiev: Naukova dumka. [Кондратюк, Е.Н., Бурда, Р.И., Остапко, В.М., 1985 – Конспект флоры Юго-Востока Украины. 272 с. Киев: Наук. думка]. (In Russian).
- Kondratyuk, E.N., Tarabrin, E.N., Baklanov, V.I., Burda, R.I., Kharkhota, A.I. (1980). *Promyshlennaya botanika*. 257 s. Kiev: Naukova dumka. [Кондратюк, Е.Н., Тарабрин, Е.Н., Бакланов, В.И., Бурда, Р.И., Хархота, А.И. Промышленная ботаника. 257 с. Киев: Наук. думка]. (In Russian).
- Konohray, V.A. (2008). *Adventywna flora Kremenchutskoho vodoskhovyshcha. Visn. Cherkas. nats. un-tu. Ser. Biol. nauky*, 128, 58–67. [Конограй, В.А., 2008 – Адвентивна флора Кременчуцького водосховища. *Вісн. Черкас. нац. ун-ту. Сер. Біол. науки*, 128, 58–67]. (In Ukrainian).
- Kostina, V.P., Bagrikova, N.A. (2010). Kostina, V.P., Bagrikova, N.A., 2010 – Annotirovannyi spisok vysshikh rasteniy Krymskogo prirodnogo zapovednika. *Nauch. zap. prirod. zapov. "Mys Martyan"*, 1, 61–142 [Костина, В.П., Багрикова, Н.А., 2010 – Аннотированный список высших растений Крымского природного заповедника. *Науч. зап. природ. запов. "Мыс Мартьян"*, 1, 61–142.]. (In Russian).
- Kotov, M.I. (1921). *Do pytannya pro te, yak poshyruyutsya teper roslyny na Ukraini zanosamy. Ukr. Bot. J.* 1(1–2), 23–25. [Котов, М.И., 1921 – До питання про те, як поширюються тепер рослини на Україні заносами. *Укр. ботан. журн.* 1(1–2), 23–25]. (In Ukrainian).
- Kotov, M.I. (1923). *Yak vplynula hromadyanska viyna na poshyrennya roslyn ta tvaryn na Ukraini ta v Rosii. Znannya* (Kharkiv), 2, 24–26. [Котов, М.И., 1923 – Як вплинула громадянська війна на поширення рослин та тварин на Україні та в Росії. *Знання* (Харків), 2, 24–26]. (In Ukrainian).
- Kotov, M.I. (1926). *K voprosu o pouyavlenii na Ukraine zanosami novykh rasteniy i o rasshirenii arealov ikh rasprostraneniya. In: Dnevnik syezda botan. (Moskva, yanvar 1926 g.)*. S. 99–100. [Котов, М.И., К вопросу о появлении на Украине заносами новых растений и о расширении ареалов их распространения. In: *Dnevnik syezda botan. (Moskva, yanvar 1926 g.)*. S. 99–100. (In Russian).

- заносами новых растений и о расширении ареалов их распространения/ В кн. Дневник съезда ботан. (Москва, янв. 1926 г.). С. 99–100]. (In Russian).
- Kotov, M.I. (1927a). Neografične doslidzhennya roslynnosti na terytorii zaliznyts Ukrainy. *Trudy silsko-hospodarskoyi botaniky*, 1(2), 133–146. [Котов, М.І., 1927а – Географічне дослідження рослинності на території залізниць України. *Тр. с.-г. ботан.*, 1(2), 133–146]. (In Ukrainian).
- Kotov, M.I. (1927b). O rasprostranenii na Ukraine *Cyclachaena xanthiifolia* Trev. *Zhurn. Rus. botan. obschestva*, 12(1–2), 197–198. [Котов, М.І., 1927b – О распространении на Украине *Cyclachaena xanthiifolia* Trev. *Журн. Рус. ботан. о-ва*. 12(1–2): 197–198]. (In Russian).
- Kotov, M.I. (1928). Adventyvna roslynnist' na Ukraini. *Visnyk pryrodoznavstva*, 5–6, 266–274. [Котов, М.І., 1928 – Адвентивна рослинність на Україні. *Вісн. природознав.*, 5–6, 266–274]. (In Ukrainian).
- Kotov, M.I. (1929). Zayshli abo adventyvni roslyny na Ukraini. *Znannia* (Kharkiv), 14, 14–16 [Котов, М.І., 1929 – Зайшли або адвентивні рослини на Україні. *Знання* (Харків), 14, 14–16]. (In Ukrainian).
- Kotov, M.I. (1934). Pro poshyrennya novykh adventyvnykh roslyn na Ukraini. *J. Inst. Bot., All Ukr. Acad. Sci.* 3(11): 99–101 [Котов, М.І., 1934 – Про поширення нових адвентивних рослин на Україні. *Журн. Ін-ту ботан. ВУАН*. 3(11), 99–101]. (In Ukrainian).
- Kotov, M.I. (1949). Adventyvni roslyny v URSR. *Bot. J. Acad. Sci. Ucr.SSR*, 6(1), 74–78 [Котов, М.І., 1949 – Адвентивні рослини в УРСР. *Ботан. журн. АН УРСР*. 6(1), 74–78]. (In Ukrainian).
- Kozhevnikova, S.K. (1970). *Adventivnye rasteniya Kryma: avtoref. diss. ... kand. biol. nauk*. 21 s. Voronezh. [Кожевникова С.К., 1970 – Адвентивные растения Крыма: автореф. дисс. ... канд. биол. наук. 21 с. Воронеж]. (In Russian).
- Kozhevnikova, S.K., Rubtsov, N.I. (1971). Opyt bioekologicheskogo i geograficheskogo analiza adventivnoy flory Kryma. *Trudy Gosudarstvennogo Nikitskoho sada*, 44, 5–93 [Кожевникова С.К., Рубцов, Н.И., 1971 – Опыт биоэкологического и географического анализа адвентивной флоры Крыма. *Тр. Гос. Никит. ботан. сада*, 44, 5–93]. (In Russian).
- Kucher, O.A. (2016). Adventivnaya fraktsiya flory Starobelskoy zlakovo-lugovoy stepi. *Fitoraznoobraznye Vostochnoy Evropy*, 10(2), 115–144 [Кучер, О.А., 2016 – Адвентивная фракция флоры Старобельской злаково-луговой степи. *Фиторазнообразие Вост. Европы*, 10(2), 115–144]. (In Russian).
- Kucher, O.A. (2017). *Adventivna fraktsiya flory Starobilskoho zlakovo-luchnoho stepu ta yiyi invazyyni potentsial: avtoref. dys. ... kand. biol. nauk*. 20 s. Kyiv [Кучер, О.О., 2017 – Адвентивна фракція флори Старобільського злаково-лучного степу та її інвазійний потенціал: автореф. дис. ... канд. біол. наук. 20 с. Київ]. (In Ukrainian).
- Kucher, O.O. (2012). Invazyyni vyd *Grindelia squarrosa* (Pursh) Dunal. na terytorii skhodu Ukrainy. In *Dynamika bioriznomanitty: Zbirnyk nauk*. pr. S. 96–99. Luhansk: Vyd-vo DZ LNU imeni Tarasa Shevchenka [Кучер, О.О., 2012 – Інвазійний вид *Grindelia squarrosa* (Pursh) Dunal. на території сходу України/ В кн.: Динаміка біорізноманіття: Зб. наук. пр. С. 96–99. Луганськ: Вид-во ДЗ ЛНУ імені Тараса Шевченка]. (In Ukrainian).

- Kucher, O.O. (2015). Transformer species in the flora of the Starobilsk grass-meadow Steppe (Ukraine). *Biodiv. Res. Conserv.* 40, 49–59.
- Kucherevskiy, V.V. (2003). *Konspekt flory Pravoberezhnoho stepovoho Prydniprov'ya*. 292 s. Dnipropetrovsk: Prospekt. [Кучеревський, В.В., 2003 – Конспект флори Правобережного степового Придніпров'я. 292 с. Дніпропетровськ: Проспект]. (In Ukrainian).
- Kucherevskiy, V.V., Shol, H.N. (2009). *Konspekt urbanoflory Kryvoho Rohu*. 71 s. Kryvyi Rih: Vydavnychiy dim [Кучеревський, В.В., Шоль, Г.Н., 2009 – Конспект урбанофлори Кривого Рогу. 71 с. Кривий Ріг: Видавн. дім]. (In Ukrainian).
- Kuzmenko, A.A. (1927). Tatarska hrechka *Fagopyrum tataricum* Gaertn. *Trudy silskohospodarskoyi botaniky*, 1(2), 88–97 [Кузьменко, А.А., 1927 – Татарська гречка *Fagopyrum tataricum* Gaertn. *Тр. с.-г. ботан.*, 1(2), 88–97]. (In Ukrainian).
- Ledebour, C. F. (1842). *Flora Rossica*. Vol. 1. 790 p. Stuttgartiae. (In Latin).
- Lukash, O.V. (2008). *Flora sudynnykh Roslyn Skhidnoho Polissya: istoriya doslidzhennya, konspekt*. 436 s. Kyiv: Fitosotsiotsentr. [Лукаш, О.В., 2008 – Флора судинних рослин Східного Полісся: історія дослідження, конспект. 436 с. Київ: Фітосоціоцентр.]. (In Ukrainian).
- Lukash, O.V. (2009). Adventyziatsiya flory sudynnykh roslyn Skhidnoho Polissya. *Ukr. Bot. J.* 66(4), 507–517. [Лукаш, О.В., 2009 – Адвентизація флори судинних рослин Східного Полісся. *Укр. ботан. журн.* 66(4), 507–517]. (In Ukrainian).
- Lytvynenko, O.I., Vynokurov, Ye.S. (2003–2004). *Amorpha fruticosa* L. – ekspansyivniy invazyivnyi vyd flory Ukrainy (polivariantnist ontogenezu ta zhyttyevist populyatsiy). *Visnyk natsionalnoho naukovoprivodnychoho muzeiu (Kyiv)*. Ser. Botan. Ch.1(2–3): 190–195. [Литвиненко, О.І., Винокуров, Є.С., 2003–2004 – *Amorpha fruticosa* L. – експансивний інвазійний вид флори України (поліваріантність онтогенезу та життєвість популяцій). *Вісн. нац. наук.-привод. музею (Київ)*. Сер. Ботан. Ч.1(2–3): 190–195]. (In Ukrainian).
- Manual of vascular plants of Ukraine (1950, 1965, 1987). [Визначник рослин України. 1950; 1965; 1987]. (In Ukrainian, and Russian).
- Mariyushkina, V.Ya. (2001). Varianty fitotsenotichnoho kontrolyu ambrozii polynolystoi. 1. *Pytannya bioindykatsii ta ekolohii*, 6(3), S. 3–18. [Мар'юшкіна, В.Я., 2001 – Варіанти фітоценотичного контролю амброзії полинолістої. 1. *Питання біоіндикації та екології*, 6(3), С. 3–18]. (In Ukrainian).
- Mariyushkina, V.Ya. (2002a). Varianty fitotsenotichnoho kontrolyu ambrozii polynolystoi. 2. *Pytannya bioindykatsii ta ekolohii*. 7(1), 10–21 [Мар'юшкіна, В.Я., 2002 – Варіанти фітоценотичного контролю амброзії полинолістої. 2. *Питання біоіндикації та екол.*, 7 (1), 10–21]. (In Ukrainian).
- Mariyushkina, V.Ya. (2002b). Varianty fitotsenotichnoho kontrolyu ambrozii polynolystoi. 3. *Pytannya bioindykatsii ta ekolohii*. 7(2–3), 39–56 [Мар'юшкіна, В.Я., 2002 – Варіанти фітоценотичного контролю амброзії полинолістої. 2. *Питання біоіндикації та екол.*, 7 (2–3), 39–56]. (In Ukrainian).
- Mariyushkina, V.Ya. (2003). *Demekolohiya invazyivnykh roslyn v ahroekosystemakh ta shlyakhy optymizatsii antropizovanykh ekosystem: avtoref. dys. ... dokt. s.-h. nauk*. 35 s. Kyiv [Мар'юшкіна, В.Я., 2003 – Демекологія інвазійних рослин в

- агроєкосистемах та шляхи оптимізації антропоізованих екосистем: автореф. дис. ... докт. с.-г. наук. – 35 с. Київ]. (In Ukrainian).
- Mariyushkina, V.Ya. (2006). *Ambroziya polynolistna: metody obstezhennya i kontrolyu: metodychni rekomendatsii*. 56 s. Kyiv: Vyd-vo Instytutu zakhystu roslyn UAAN [Мар'юшкіна, В.Я., 2006 – Амброзія полинолиста: методи обстеження і контролю: методичні рекомендації. 56 с. Київ: Вид-во Ін-ту захис. росл. УААН]. (In Ukrainian).
- Mariyushkina, V.Ya., Hrytsenko, V.V. (2000). Perspektyvy fitotsenotychnoho kontrolyu *Solidago canadensis* L. *Dopovidi NAN Ukrainy*, 8, 158–162. [Мар'юшкіна, В.Я., Гриценко, В.В., 2000 – Перспективи фітоценотичного контролю *Solidago canadensis* L. *Доп. НАН України*, 8, 158–162]. (In Ukrainian).
- Mariyushkina, V.Ya. (1986). *Ambroziya polynolistnaya i osnovy biologicheskoy borby s ney*. 117 s. Kiev: Nauk. dumka. [Мар'юшкіна, В.Я., 1986 – Амброзія полыннолистная и основы биологической борьбы с ней. 117 с. Киев: Наук. думка]. (In Russian).
- Melnychuk, S.S., Trokhymenko, H.H. (2014). Adventywna fraktsiya flory Natsionalnoho pryrodnoho parku “Biloberezhzhya Svyatoslava”. *Naukovyi visnyk Mykolaivskoho derzhavnoho universytetu im. V.O. Sukhomlynskoho*. Ser. Biol. nauky., 6.3 (113): 68–73. [Мельничук, С.С., Трохименко, Г.Г., 2014 – Адвентивна фракція флори Національного природного парку “Білобережжя Святослава”. *Наук. вісн. Миколаїв. держ. ун-ту ім. В.О. Сухомлинського*. Сер. Біол. Науки., 6.3 (113), 68–73]. (In Ukrainian).
- Melnyk, R.P. (2001). *Urbanoflora Mykolayeva: avtoref. dys. ... kand. biol. nauk*. 19 s. Yalta. [Мельник, Р.П., 2001 – Урбанофлора Миколаєва: автореф. ... канд. біол. наук, 19 с. Ялта]. (In Ukrainian).
- Montrezor, V.V. (1886, 1887, 1888, 1890a, 1890b). Obozrenie rasteniy, vkhodyashchikh v sostav flory guberniy Kievskogo uchebnogo okruga: Kievskoy, Podolskoy, Volynskoy, Chernigovskoy i Poltavskoy. *Zapiski Kievskogo obschatva estestvoispytateley*, 8(1), 1–144; 8(2), 185–288; 9(1–2), 119–198; 10(3), 457–546; 10(4), 1–90. [Монтрезор, В.В., 1886 – Обзорение растений, входящих в состав флоры губерний Киевского учебного округа: Киевской, Подольской, Волынской, Черниговской и Полтавской. *Зап. Киев. о-ва естествоисп.* 8(1), 1–144; 8(2), 185–288; 9(1–2), 119–198; 10(3), 457–546; 10(4), 1–90]. (In Russian).
- Mosyakin, S.L. (1988). Rid *Bidens* L. (Asteraceae) u flori URSR. *Ukr. Bot. J.*, 45(6), 63–64 [Мосякін, С.Л., 1988 – Рід *Bidens* L. (Asteraceae) у флорі УРСР. *Укр. ботан. журн.*, 45(6), 63–64]. (In Ukrainian).
- Mosyakin, S.L. (1989). Vudy rodu *Setaria* Beauv. (Poaceae) u flori URSR. *Ukr. Bot. J.*, 46(5), 33–35 [Мосякін, С.Л., 1989 – Види роду *Setaria* Beauv. (Poaceae) у флорі УРСР. *Укр. ботан. журн.*, 46(5), 33–35]. (In Ukrainian).
- Mosyakin, S.L. (1992). The genus *Puccinellia* Parl. (Poaceae) in the Ukraine. *Ukr. Bot. J.*, 49(1), 11–14.
- Mosyakin, S.L. (1995a). Rid *Cenchrus* L. (Poaceae) v Ukraini: ohlyad nomenklatury, systematyky ta suchasnoho poshyrennya. *Ukr. Bot. J.*, 52(1), 120–126. [Мосякін, С.Л., 1995 – Рід *Cenchrus* L. (Poaceae) в Україні: огляд

- номенклатури, систематики та сучасного поширення. *Укр. ботан. журн.*, 52(1), 120–126]. (In Ukrainian).
- Mosyakin, S.L. (1995b). Ohlyad rodu *Amaranthus* L. (Amaranthaceae) v Ukraini. *Ukr. Bot. J.*, 52(2), 65–74 [Мосякін, С.Л., 1995б – Огляд роду *Amaranthus* L. (Amaranthaceae) в Україні. *Укр. ботан. журн.*, 52(2), 65–74]. (In Ukrainian).
- Mosyakin, S.L., Korniyenko, O.M. (2006). Nomenklatura kultyvovanykh ta zdychavilykh v Ukraini pivnichnoamerykanskykh “aystr” z tochky zoru delimitatsii rodiv u trybi Astereae (Asteraceae). *Ukr. Bot. J.*, 63(2), 159–165. [Мосякін, С.Л., Корнієнко, О.М., 2006 – Номенклатура культивованих та здичавілих в Україні північноамериканських “айстр” з точки зору делімітації родів у трибі Astereae (Asteraceae). *Укр. ботан. журн.*, 63(2), 159–165]. (In Ukrainian).
- Mosyakin, S.L., Yavorska, O.G. (2002). The nonnative flora of the Kyiv (Kiev) Urban Area, Ukraine: A checklist and brief analysis. *Urban Habitats*, 1(1) (Spec. iss.: Urban floras). Available at: http://www.urbanhabitats.org/v01n01/nonnativekiev_full.html
- Moysiienko, I.I. (1999). *Urbanoflora Khersona: avtoref. dys. ... kand. biol. nauk.* 19 s. Yalta. [Мойсієнко, І.І., 1999 – Урбанофлора Херсона: автореф. дис. ... канд. біол. наук. 19 с. Ялта]. (In Ukrainian).
- Moysiienko, I.I. (2011). *The Flora of the Pivnichne Prychornomor'ia (Structural Analysis, Synantropization, Conservation): avtoref. dys. ... dokt. biol. nauk.* 35 s. Kyiv. [Мойсієнко, І.І., 2011 – Флора Північного Причорномор'я (структурний аналіз, синантропізація, охорона): автореф. дис. ... докт. біол. наук. 35 с. Київ]. (In Ukrainian).
- Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, Th., Čárni, A., Šumberová, K., Willner, W., Dengler, J., García, R. G., Chytrý, M., Hájek, M., Di Pietro, R., Iakushenko, D., Pallas, J., Daniels, F. J.A., Bergmeier, E., Guerra, A. S., Ermakov, N., Valachovič, M., Schaminée, J. H.J., Lysenko, T., Didukh, Y.P., Pignatti, S., Rodwell, J.S., Capelo, J., Weber, H.E., Solomeshch, A., Dimopoulos, P., Aguiar, C., Hennekens, S.M., Tichý, L. (2016). Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Appl. Veg. Science*, 19 (Suppl. 1), 3–264. Doi: 10.1111/avsc.12257
- Nechytaylo, V.A., Pohrebennyk, V.P., Hryshchenko, V.V. (2002). *Sudynni roslyny Kanivskoho zapovidnyka i okolyts*, 226 s. Kyiv: Fitosotsiotsentr. [Нечитайло, В.А., Погребенник, В.П., Грищенко, В.В., 2002 – Судинні рослини Канівського заповідника і околиць, 226 с. Київ: Фітосоціоцентр]. (In Ukrainian).
- Oitsius', L.V. (2011). *Adventywni vydy flory Volynskoho Polissia: avtoref. dys. ... kand. biol. nauk*, 24 s. Kyiv [Ойцюсь, Л.В., 2011 – Адвентивні види флори Волинського Полісся: автореф. дис. ... канд. біол. наук. 24 с. Київ]. (In Ukrainian).
- Oliyar, H.I. (2002). Konspekt flory pryrodnoho zapovidnyka “Medobory”. *Naukovi zapysky Ternopil'skoho derzhavnoho pedahohichnoho universytetu*, Ser. Biol., 2(17), 18–25. [Оліяр, Г.І., 2002 – Конспект флори природного заповідника

- “Медобори”. *Наук. зап. Тернопіль. держ. пед. ун-ту. Сер. Біол.*, 2(17), 18–25]. (In Ukrainian).
- Orlov, A.A. (red.) (2013). *Fitoriznomanittia Poliskoho pryrodnoho zapovidnyka: vodorosti, mokhopodibni, sudynni roslyny*. 256 s. Kyiv: NVP-Interservis. [Фіторізноманіття Поліського природного заповідника: водорості, мохоподібні, судинні рослини / Ред. А.А., Орлов. Київ: НВП-Інтерсервіс]. (In Ukrainian).
- Ostapko, V.M., Boyko, A.V., Mosyakin, S.L. (2010). *Sosudistye rasteniya Yugo-Vostoka Ukrainy*. 247 s. Donetsk: Noulidzh. [Остапко, В.М., Бойко, А.В., Мосякин, С.Л., 2010 – Сосудистые растения Юго-Востока Украины. 247 с. Донецк: Ноулидж.]. (In Russian).
- Ostapko, V.M., Boyko, A.V., Mulenkova, O.G. (2009). Adventivnaya fraktsiya flory Yugo-Vostoka Ukrainy. *Promyshlennaya botanika: Sbornik nauchnykh trudov*, 9, 32–47 [Остапко, В.М., Бойко, А.В., Муленкова, О.Г., 2009 – Адвентивная фракция флоры Юго-Востока Украины. *Промыш. ботан.*: Сб. науч. тр., 9, 32–47]. (In Russian).
- Paczoski, I.K. (1897, 1899, 1900). Flora Polesya i prilezhashchikh mestnostey. *Trudy S.-Peterburgskogo obschestva estestvoispytateley*, 27(2), 1-103 p., 29 (3): 1-113, 30 (3), 1-259. [Пачоский, И.К., 1897, 1899, 1900 – Флора Полесья и прилежащих местностей, *Тр. С.-Петербур. о-ва естествоисп.*, 27(2), 1-103 p., 29 (3), 1-113, 30 (3), 1-259]. (In Russian).
- Paczoski, I.K. (1911). O sorno-polevoy rastitelnosti Khersonskoy gubernii. *Trudy byuro po prikladnoy botanike*, 4(3), 71–146 [Пачоский, И.К., 1911 – О сорно-полевой растительности Херсонской губернии. *Тр. бюро по прикл. ботан.* 4(3), 71–146]. (In Russian).
- Panchenko, S.M. (2005a). *Flora Natsionalnoho pryrodnoho parku «Desnyansko-Starohutskiy» ta problemy okhorony fitoriznomanittya Novhorod-Siverskoho Polissya*. 170 s. Sumy: Universytetska knyha [Панченко, С.М., 2005а – Флора Национального природного парка “Деснянсько-Старогутський” та проблеми охорони фіторізноманіття Новгород-Сіверського Полісся. 170 с. Суми: Університетська книга]. (In Ukrainian).
- Panchenko, S.M. (2005b). Invaziyna spromozhnist pivnichno-amerykanskooho vydu *Conyza canadensis* (L.) Cronq. v NPP “Desnyansko-Starohutskiy”. *Ukr. Bot. J.*, 62(4), 558-564 [Панченко, С.М., 2005б – Інвазійна спроможність північно-американського виду *Conyza canadensis* (L.) Cronq. в НПП “Деснянсько-Старогутський”. *Укр. ботан. журн.*, 62(4), 558-564]. (In Ukrainian).
- Petryk, S.P. (2006). Pro vdoskonalennya systemy fito-sanitarnoho kontrolyu v Ukraini. *Ahrarnyi visnyk Prychornomia: zbirnyk naukovykh prats*, 29, 269–272 [Петрик, С.П., 2006 – Про вдосконалення системи фіто-санітарного контролю в Україні. *Аграр. вісн. Причорн.*: Зб. наук. пр., 29: 269–272]. (In Ukrainian).
- Prokudin, Yu.M., Vovk, A.G., Petrova, O.A., Ermolenko, E.D., Vernichenko, Yu.V. (1977). *Zlaki Ukrainy*. 518 s. Kiev: Nauk. dumka. [Прокудин, Ю.М., Вовк, А.Г., Петрова, О.А., Ермоленко, Е.Д., Верниченко, Ю.В., 1977 – Злаки Украины. 518 с. Киев: Наук. думка]. (In Russian).
- Protopopova, V., Shevera M.V., Anischenko, I.M., Terentyeva, N.H. (2010b). Analiz kenofitiv urbanoflor riznykh botaniko-geografichnykh zon Ukrainy iz zastosuvanniam metodiv matematychnoyi statystyky. *Ukr. Bot. J.*, 67(4), 536–

- 546 [Протопопова, В.В., Шевера, М.В., Аніщенко, І.М., Терентьєва, Н.Г., 2010 – Аналіз кенофітів урбанофлор різних ботаніко-географічних зон України із застосуванням методів математичної статистики. *Укр. ботан. журн.* 67(4), 536–546. (In Ukrainian).
- Protopopova, V., Shevera, M. (2002b). *A preliminary checklist of the urban flora of Uzhgorod*. 68 p. Kyiv: Phytosociocentre.
- Protopopova, V., Shevera, M. (2008a). Participation of archaeophytes in urban floras in different botanical and geographical zones of Ukraine: a preliminary assessment, *Thaiszia. – J. Bot.* (Košice), 18, Suppl. 1, 89–104.
- Protopopova, V., Shevera, M. (2008b). Participation of alien species in urban floras in different botanical and geographical zones of Ukraine: a preliminary assessment, *Biodiv. Res. Conserv.*, 11–12, 9–16.
- Protopopova, V., Shevera, M.V. Anischenko I.M., Terentyeva N.H. (2013). Porivnialna kharakterystyka archeofitiv urbanoflor riznykh botaniko-geografichnykh zon Ukrainy. *Ukr. Bot. J.*, 70(2), 158–163 [Протопопова, В.В., Шевера, М.В., Аніщенко, І.М., Терентьєва, Н.Г., 2013 – Порівняльна характеристика археофітів урбанофлор різних ботаніко-географічних зон України. *Укр. ботан. журн.*, 70(2), 158–163. (In Ukrainian).
- Protopopova, V.V. (1964). Novi dani pro systematychnyi sklad rodu netreba (*Xanthium* L.) na Ukraini. *Ukr. Bot. J.*, 21(4), 78–84. [Протопопова, В.В., 1964 – Нові дані про систематичний склад роду нетреба (*Xanthium* L.) на Україні. *Укр. ботан. журн.*, 21(4), 78–84. (In Ukrainian).
- Protopopova, V.V. (1965). Adventyvni roslyny Lisostepu ta Stepu Ukrainy. *Ukr. Bot. J.*, 22 (3), 38–44. [Протопопова, В.В., 1965 – Адвентивні рослини Лісостепу та Степу України. *Укр. ботан. журн.*, 22(3), 38–44]. (In Ukrainian).
- Protopopova, V.V. (1973). *Adventyvni roslyny Lisostepu i Stepu Ukrainy*, 188 s. Kyiv: Naukova dumka. [Протопопова, В.В. 1973 – Адвентивні рослини Лісостепу і Степу України. 188 с. Київ: Наук. думка]. (In Ukrainian).
- Protopopova, V.V. (1974). Do vydivoho skladu kvasenyts (*Oxalis* L.) na Ukraini. *Ukr. Bot. J.*, 31(1), 110–113 [Протопопова, В.В., – 1974 – До видового складу квасениць (*Oxalis* L.) на Україні. *Укр. ботан. журн.*, 31(1), 110–113]. (In Ukrainian).
- Protopopova, V.V. (1991). *Synantropnaya flora Ukrainy i puti yeyo rozvitiya*, 204 s. Kiev: Nauk. dumka. [Протопопова, В.В., 1991 – Синантропная флора Украины и пути ее развития. 204 с. Киев: Наук. думка]. (In Russian).
- Protopopova, V.V., Mosyakin, S.L., Shevera, M.V. (2002a). *Fitoinvazii v Ukraini yak zahroza bioriznomanittyu: suchasnyi stan i zavdannya na maybutnye*. 32 s. Kyiv: Inst. botaniky NAN Ukrainy. – [Протопопова, В.В., Мосякін, С.Л., Шевера, М.В., 2002 – Фітоінвазії в Україні як загроза біорізноманіттю: сучасний стан і завдання на майбутнє. 32 с. Київ: Ін-т ботаніки НАН України]. (In Ukrainian).
- Protopopova, V.V., Mosyakin, S.L., Shevera, M.V. (2003). Vplyv neaboryhennykh vydiv roslyn na biotu Ukrainy. In: *Otsinka i napryamky zmeshhennya zahroz bioriznomanittyu / Red. O. Dudkin*. Ss. 129–155; 358; 364–374, Kyiv: Khimdzhest. [Протопопова, В.В., Мосякін, С.Л., Шевера, М.В., 2003 – Вплив неаборигенних видів рослин на біоту України. В кн.: Оцінка і напрямки зменшення загроз біорізноманіття / Ред. О. Дудкін. С. 129–155; 358; 364–374. Київ: Хімджест, С. 129–155; 358; 364–374]. (In Ukrainian).

- Protopopova, V.V., Shevera, M.V. (2014b). Ergasiophytes of the Ukrainian flora. *Biodiv. Res. Conserv.*, 35, 31–46.
- Protopopova, V.V., Shevera, M.V., Bagrikova, N.O., Ryff, L.E. (2012). Vyd-transformery u flori Pivdennoho bereha Krymu. *Ukr. Bot. J.*, (69)1, 54–68. [Протопопова, В.В., Шевера, М.В., Багрікова, Н.О., Рифф, Л.Е., 2012 – Види-трансформери у флорі Південного берега Криму. *Укр. ботан. журн.*, (69)1, 54–68]. (In Ukrainian).
- Protopopova, V.V., Shevera, M.V., Chorney, I.I., Tokaryuk, A.I., Budzhak, V.V., Korzhan, K.V. (2010a). Vyd-transformery u flori Bukovynskoho Peredkarpattya. *Ukr. Bot. J.*, (67)6, 852–863. [Протопопова, В.В., Шевера, М.В., Чорней, І.І., Токарюк, А.І., Буджак, В.В., Коржан, К.В., 2010 – Види-трансформери у флорі Буковинського Передкарпаття. *Укр. ботан. журн.*, (67)6, 852–863]. (In Ukrainian).
- Protopopova, V.V., Shevera, M.V., Fedoronchuk, M.M., Shevchuk, V.L. (2014a). Vyd-transformery u flori Srednyoho Prydniprovia. *Ukr. Bot. J.*, 71(5), 563–572. [Протопопова, В.В., Шевера, М.В., Федорончук, М.М., Шевчук, В.Л., 2014 – Види-трансформери у флорі Середнього Придніпров'я, *Укр. ботан. журн.*, 71(5), 563–572]. (In Ukrainian).
- Protopopova, V.V., Shevera, M.V., Mosyakin, S.L. (2006). Deliberate and unintentional introduction of invasive weeds: a case study of the alien flora of Ukraine. *Euphytica*, 148, 17–33.
- Protopopova, V.V., Shevera, M.V., Mosyakin, S.L., Solomakha, V.A., Solomakha, T.D., Vasylyeva, T.V., Petryk, S.P. (2009). Vyd-transformery u flori Pivnichnoho Prychornomor'ya. *Ukr. Bot. J.*, 66(6), 770–782. [Протопопова, В.В., Шевера, М.В., Мосякін, С.Л., Соломаха, В.А., Соломаха, Т.Д., Васильєва, Т.В., Петрик, С.П., 2009 – Види-трансформери у флорі Північного Причорномор'я. *Укр. ботан. журн.*, 66(6), 770–782]. (In Ukrainian).
- Protopopova, V.V., Shevera, M.V., Orlov, O.O., Panchenko, S.M. (2015). The transformer species of the Ukrainian Polissya. *Biodiv. Res. Conserv.*, 39, 7–18.
- Protopopova, V.V., Tyukh, YuYu., Shevera, M.V. (1999). Kharakterystyka adventyvnoyi fraktsiyi flory Natsionalnoho pryrodnoho parku "Synevyr". In: Problemy ekologichnoyi stabilnosti Skhidnykh Karpat: Mizhnarodna naukovopraktychna konferentsia, prysviachena 10-richchiu stvorennia NPP "Synevyr": Materialy konferencii (Synevyr, 24–27 chervnia 1999 r.). S. 157–158. [Протопопова, В.В., Тюх, Ю.Ю., Шевера, М.В., 1999 – Характеристика адвентивної фракції флори Національного природного парку "Синевир". В кн.: Проблеми екологічної стабільності Східних Карпат: Міжн. наук.-практ. конф., присвяч. 10-річчю створ. НПП "Синевир": Мат. конф. (Синевир, 24–27 черв. 1999 р.), С. 157–158]. (In Ukrainian).
- Prots, B. (2015). Invasion success of alien plant species in the Carpathian region of Ukraine. *Visnyk Chernivtsi univ. Biol. syst.*, 5(1), 116–121.
- Prots, B., Drescher, A., Vykhor, B. (2011). Invasion ecology of Green Ash *Fraxinus pennsylvanica* Marshall in the Transcarpathia (Ukraine). *Visnyk Chernivtsi univ. Biol. syst.*, 3(3), 269–276.

- Rogovich, A.S. (1855). Obozrenie sosudistyykh I polusosudistyykh rasteniy, vkhodyashchikh v sostav flory guberniy: Kievskoy, Chernigovskoy I Poltavskoy. *Trudy Komissii pry universitete sv. Vladymyra dlya opisaniya huberniy Kievskogo uchebnogo okruga*, 3, 1–147 [Рогович, А.С., 1855 – Обзорение сосудистых и полусосудистых растений, входящих в состав флоры губерний: Киевской, Черниговской и Полтавской. *Тр. Комис. для описания губерний Киевского учебного округа*, 3, 1–147]. (In Russian).
- Rogovich, A.S. (1869). *Obozrenie semennykh i vysshikh sporovykh rasteniy, vkhodyashchikh v sostav flory guberniy Kievskogo uchebnogo okruga: Volynskoy, Podolskoy, Kievskoy, Chernigovskoy i Poltavskoy*. 308 s. Kiev: Kiev Universitet. [Рогович, А.С., 1869 – Обзорение семенных и высших споровых растений, входящих в состав флоры губерний Киевского учебного округа: Волынской, Подольской, Киевской, Черниговской и Полтавской, 308. Киев: Изд. Киев. ун-та. (In Russian).
- Rostański, K., Dzhus, M., Gudžinskas, Z., Rostański A., Shevera, M.V., Šulcs, V., Tokhtar, V. (2004). *The genus Oenothera L. in Eastern Europe*. 133 p. Krakow: W. Szafer Institute of Botany, PAN.
- Shevchyk V.L., Senchylo O.O. (2009). Adventyvnа flora zaplavy Kanivskoho pryrodnoho zapovidnyka. *Chornomorskyi Bot. J.*, 5(4), 563–570. [Шевчик В.Л., Сенчило О.О., 2009 – Адвентивна флора заплави Канівського природного заповідника, *Чорном. ботан. журн.*, 5(4), 563–570]. (In Ukrainian).
- Shevera, M.V. (2005). Urbanoflory Ukrainy: sovremennoe sostoyanie izuchennosti. V kn.: *Izuchenie flory Vostochnoy Evropy: dostizheniya i perspektivy*. Tez. dokl. Mezhd. konf. (Sankt-Peterburg, 23–28 maya 2005). S. 101. Moskva, Sankt-Peterburg: tov-vo nauchn. Izdaniy KMK. [Шевера, М.В., 2005 – Урбанофлоры Украины: современное состояние изученности. В кн.: Изучение флоры Восточной Европы: достижения и перспективы. Тез. докл. Межд. конф. (Санкт-Петербург, 23–28 мая 2005). С. 101. Москва, Санкт-Петербург: тов-во научн. Изданий КМК]. (In Russian).
- Shmalhausen, I.F. (1886). *Flora Yugo-Zapadnoy Rossii*, 783. Kiev. [Шмальгаузен, И.Ф., 1886 – Флора Юго-Западной России, 783. Киев. (In Russian).
- Shol, H. (2004). Flora Kryvoho Rohu: suchasnyi stan ta sozologichni aspekty. *Visn. Lviv. Univ., Ser. Biol.*, 36, 63-69 [Шоль, Г., 2004 – Флора Кривого Рогу: сучасний стан та созологічні аспекти. *Вісн. Львів. ун-ту. Сер. Біол.*, 36, 63-69]. (In Ukrainian).
- Taliev, V.I. (1900). Flora Kryma i rol' cheloveka v yeye razvitii. *Trudy obschestva ispytateley prirody pri Kharkovskom universitete*, 35, 220–221. [Талиев, В.И., 1900. – Флора Крыма и роль человека в ее развитии, *Тр. о-ва испыт. природы при Харьков. ун-те*, 35, 220–221. (In Russian).
- Taliev, V.I. (1902). Chelovek kak botaniko-geograficheslyi factor. *Nauchnoye obozreniye*, 11, 42–61. [Талиев, В.И., 1902 – Человек как ботанико-географический фактор, *Науч. обозрение*, 11, 42–61. (In Russian).
- Tarasov, V.V. (2005). *Flora Dnipropetrovskoi ta Zaporizkoi oblastey. Sudynni roslyny*. 276 s. Dnipropetrovsk: Vyd-vo Dnipropetrosk Nats. Univ. [Тарасов, В.В., 2005 – Флора Дніпропетровської та Запорізької областей. Судинні рослини. 276 с. Дніпропетровськ: Вид-во Дніпропетровськ. нац. ун-ту]. (In Ukrainian).

- Tarasov, V.V. (2012). *Flora Dnipropetrovskoi ta Zaporizkoi oblastey*. 296 s. Dnipropetrovsk: Lira. [Тарасов, В.В. 2012 – Флора Дніпропетровської та Запорізької областей. 296 с. Дніпропетровськ: Ліра]. (In Ukrainian).
- Tokaryuk, A.I., Chorney, I.I., Budzhak, V.V., Protoporova, V.V., Shevera, M.V., Korzhan, K.V., Volutsa, O.D. (2018). *Invaziyini roslyny u Bukovynskomu Peredkarpatti*. 200 s. Chernivtsi: Druk Art. [Токарюк, А.І., Чорней, І.І., Буджак, В.В., Протопопова, В.В., Шевера, М.В., Коржан, К.В., Волуца, О.Д., 2018 – Інвазійні рослини у Буковинському Передкарпатті. 200 с. Чернівці: Друк Арт]. (In Ukrainian).
- Tokaryuk, A.I., Chorney, I.I., Korzhan, K.V., Budzhak, V.V., Velychko, M.V., Protoporova, V.V., Shevera, M.V. (2012). The participation of invasive plants in the synanthropic plant communities in the Bukovinian Cis-Carpathians (Ukraine). *Thaiszia – J. Bot.*, 22(2), 243-254.
- Tokhtar, V.K. (2005). *Flora tekhnogennykh ekotopov i ee razvitie (na primere yugovostoka Ukrainy: avtoref. dis. ... dokt. biol. nauk*. 36 s. Kyiv. [Тохтарь, В.К., 2005 – Флоры техногенных экотопов и их развитие (на примере юго-востока Украины): автореф. дис... д-ра биол. наук. 36 с. Киев]. (In Russian).
- Tokhtar', V.K. (1993). *Adventyvna flora zaliznyts Pivdennoho-sходу Ukrainy: avroref. dys. ... kand. biol. nauk*, 18 s. Kyiv [Тохтарь, В.К., 1993 – Адвентивна флора залізниць Південного-сходу України: автореф. дис. ... канд. біол. наук. 18 с. Київ]. (In Ukrainian).
- Vasylyeva, T.V., Kovalenko, S.H. (2003). *Konspekt flory pivdennoi Bessarabii*. 250 s. Odesa: Odeskyi Nats. Un-t im. I.I. Metchnykova [Васильєва, Т.В., Коваленко, С.Г., 2003 – Конспект флори південної Бессарабії. 250 с. Одеса: Одеський нац. ун-т ім. І.І.Мечникова]. (In Ukrainian).
- Vasylyeva-Nemertsalova, T.V. (1996). *Synantropna flora pryportovykh mist Pivnichno-Zakhidnoho Prychornomor'ya ta shlyakhy yiyi rozvytku: avtoref. dys. ... kand. biol. nauk*. 21 s. Kyiv. [Васильєва-Немерцалова, Т.В., 1996 – Синантропна флора припортових міст Північно-Західного Причорномор'я та шляхи її розвитку: автореф. дис. ... канд. біол. наук. 21 с. Київ]. (In Ukrainian).
- Vykhor, B., Prots, B. (2012). Borshchivnyk Sosnovskoho (*Heracleum sosnowskyi* Manden.) na Zakarpatti: ekolohiya, poshyrennya ta vplyv na dovkillya. *Biol. studii / Studia Biologica*, 6(3), 185–196 [Вихор, Б.І., Проць, Б.Г. Борщівник Сосновського (*Heracleum sosnowskyi* Manden.) на Закарпатті: екологія, поширення та вплив на довкілля. *Біол. студії / Studia Biologica*, 6(3), 185–196]. (In Ukrainian).
- Vykhor, B., Prots, B. (2014). Invaziyini vydy roslyn Zakarpattya: ekolohichna kharakterystyka ta dynamichni tendentsii poshyrennya. *Biol. stud. – Stud. Biol.*, 8(1), 171–186 [Вихор, Б., Проць, Б., 2014 – Інвазійні види рослин Закарпаття: екологічна характеристика та динамічні тенденції поширення. *Біол. студ. – Stud. Biol.*, 8(1), 171–186]. (In Ukrainian).
- Vykhor, B.I. (2015). *Ekolohichna otsinka vplyvu invaziynykh vydiv roslyn na fitoriznomanitnya Zakarpattya: avtoref. dys. ... kand. biol. nauk*. 20 s. Kyiv. [Вихор, Б.І., 2015 – Екологічна оцінка впливу інвазійних видів рослин на фіторізноманіття Закарпаття: автореф. дис. ... канд. біол. наук. 20 с. Київ]. (In Ukrainian).

- Yanata, O.A. (1927). Bur'yany Ukrainy – vyvchennya yikh ta sposib borotby z nymy. *Trudy silsko-hospodarskoyi botaniky*, 1(2), 3–16 [Яната, О.А. 1927 – Бур'яни України – вивчення їх та спосіб боротьби з ними, *Тр. с.-г. ботан.* 1(2), 3–16]. (In Ukrainian).
- Zapałowicz, H. (1906, 1911a, 1911b). *Conspectus florum Galiciae criticus*, Т. 1, 296 s., Т. 2, 311 s., Т. 3, 246 s. Kraków: Akad. Umiejętności. (In Poland).
- Zavalova, L.V. (2008). Alien fraction of Chernihiv urban flora: analysis and checklist. *Biodiv. Res. Conserv.*, 11-12, 17–26.
- Zavalova, L.V. (2010). *A Checklist of Chernihiv urban flora* / Sci. Ed. M. Shevera. 107 p. Kyiv: Phytosociocenter, M.G. Kholodny Institute of Botany, NAS of Ukraine.
- Zavalova, L.V. (2012). *Urbanoflora Chernihova: avroref. dys. ... kand. biol. nauk.* 22 s. Kyiv. [Зав'ялова, Л.В., 2012 – Урбанофлора Чернігова: автореф. дис. ... канд. біол. наук. 22 с. Київ]. (In Ukrainian).
- Zavalova, L.V. (2017). Invazionniye vidy rasteniy vo flore zapovednykh obyektov Ukrainy. *Botanika (issledovaniye): Sbornik nauchnykh trudov* (Minsk), 46, 27–39 [Зав'ялова, Л.В., 2017 – Инвазионные виды растений во флоре заповедных объектов Украины, *Ботаника (исследования): Сб. научн. тр.* (Минск), 46, 27–39]. (In Russian).
- Zavalova, L.V. (2017). Vudy invaziynikh roslyn, nebezpechni dla pryrodnoho fitoriznomanittia obyektiv pryrodno-zapovidnoho fondu Ukrainy. *Naukovyi visnyk Chernivetskoho Natsionalnoho Universytetu. Biol. (Biologichni systemy)*, 9(1), 87–107 [Зав'ялова, Л.В., 2017 – Види інвазійних рослин, небезпечні для природного фіторізноманіття об'єктів природно-заповідного фонду України. *Наук. вісн. Чернівецьк. нац. ун-ту. Біол. (Біол. системи)*, 9(1), 87–107]. (In Ukrainian).
- Zavalova, L.V., Korniyenko, O.M. (2014). Fitozabrudnennya Dniprovskoho ekolohichnoho korydoru In: *Ekolohiya vodno-bolotnykh uhid i torfovnyshch.* P. 90–93. Kyiv: Interservis. [Зав'ялова, Л.В., Корнієнко, О.М., 2014 – Фітозабруднення Дніпровського екологічного коридору. У кн.: *Екологія водно-болотних угідь і торфовищ.* С. 90-93. Київ: Інтерсервіс]. (In Ukrainian).
- Zawadzki, A. (1835). *Enumeratio plantarum Galiziae a Bukowinae oder die in Galizien und der Bukowina wild wachsenden Pflanzen*, 200 s. Breslau. (In German)
- Zviagintseva, K.O. (2015). *An annotated checklist of the urban flora of Kharkiv* / Sci. ed. M. Shevera. 96 p. Kharkiv: V.N. Karazin Kharkiv National University.
- Zviagintseva, K.O. (2015). *Antropohenna transformatsiya flory m. Kharkova: avtoref. dis. ... kand. biol. nauk.* 19 s. Kyiv. [Звягінцева, К.О. (2015). Антропогенна трансформація флори м. Харкова: автореф. дис. ... канд. біол. наук. 19 с. Київ]. (In Ukrainian).